

Additive Toxicity Statement

2-Propenoic acid, butyl ester

DESCRIPTION

Name 2-Propenoic acid, butyl ester


IUPAC name Butyl prop-2-enoate

CAS no 141-32-2

EINECS no. 205-480-7

Molecular formula $C_7H_{12}O_2$

Structural formula



Molecular weight (g/mol) 128.2

CLP HEALTH HAZARD CLASSIFICATION

This ingredient has been classified under Regulation (EC) No 1272/2008 of the European Parliament and of the Council and is in the classification and labelling inventory.

The CLP classifications of this ingredient are not considered applicable for one, or more, of the following reasons:

- the level required to elicit a response in a toxicological study, with a relevant route of exposure, is far higher than that of its use in BAT products
- the toxicological effect identified does not relate to the route of exposure associated with its use in BAT products
- no relevant toxicological study was identified relating to the route of exposure, and there is only limited consumer exposure to this ingredient associated with its use in BAT products

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are: TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this, our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch *et al*¹¹.

In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴
- Assessment of the irritation potential of Swedish snus ingredients using the Epioral™ tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself. In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

1. Available at: [http://www.bibra-information.co.uk/supported access to our chemical toxicology database TRACE.html](http://www.bibra-information.co.uk/supported%20access%20to%20our%20chemical%20toxicology%20database%20TRACE.html)
2. Available at: <http://toxnet.nlm.nih.gov/index.html>
3. Available at: <http://ccinfoweb.ccohs.ca/rtecs/search.html>
4. Available at: <http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384>
5. Available at: <http://www.inchem.org/>
6. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
7. Available at: <http://echa.europa.eu/information-on-chemicals>
8. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>
9. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp>
10. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
11. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
12. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem. Toxicol.* 42 Suppl:S53-S83, 2004.
13. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids. *Regul. Toxicol. Pharmacol.* 72 (2):361-369, 2015.
14. B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. *Toxicology Letters* 229S:S109, 2014. (Abstract)
15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. *The Toxicologist*, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
16. World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004
17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf.
18. SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_051.pdf.

Additive Toxicity Statement

2-Propenoic acid, butyl ester



REFERENCES

- 3M Belgium BVBA/SPRL (OR1) *et al.* (2023). REACH registration dossier on butyl acrylate (CAS RN 141-32-2). Disseminated on the European Chemicals Agency (ECHA) website. Last modified 28-Mar-2023. <https://echa.europa.eu/fi/registration-dossier/-/registered-dossier/15779>
- Anon. (2017). Reproductive toxicity. 001 Key / Experimental result (cited in 3M Belgium BVBA/SPRL (OR1) *et al.*, 2023). <https://echa.europa.eu/fi/registration-dossier/-/registered-dossier/15779/7/9/2>
- Baur X (2013). A compendium of causative agents of occupational asthma. *Journal of Occupational Medicine and Toxicology* 8, 15.
- Burdock GA (2010). *Fenaroli's Handbook of Flavor Ingredients*. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.
- ChemSpider. Royal Society of Chemistry chemical structure database. Record for butyl acrylate (CAS RN 141-32-2). Accessed Oct-2023. <https://www.chemspider.com/Chemical-Structure.8514.html>
- DFG (2018). Deutsche Forschungsgemeinschaft. n-Butyl acrylate. The MAK Collection for Occupational Health and Safety 2018, Vol 3, No. 2, 455-465. <https://onlinelibrary.wiley.com/doi/pdf/10.1002/3527600418.mb14132e6218>
- EC (2008a). European Commission. List of flavourings and source materials as laid down in Part I of Annex I of Regulation (EC) No 1334/2008. <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2008:354:0034:0050:en:PDF>
- EC (2008b). European Commission. Regulation (EC) No 1334/2008 of the European Parliament and of the Council of 16 December 2008 on flavourings and certain food ingredients with flavouring properties for use in and on foods and amending Council Regulation (EEC) No 1601/91, Regulations (EC) No 2232/96 and (EC) No 110/2008 and Directive 2000/13/EC. <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=celex:32008R1334>
- ECHA (a). European Chemicals Agency. Candidate List of substances of very high concern (SVHC) for Authorisation. Accessed Oct-2023. <https://echa.europa.eu/candidate-list-table>
- ECHA (b). European Chemicals Agency. Authorisation List. Accessed Oct-2023. <https://echa.europa.eu/authorisation-list>
- ECHA (2023). European Chemicals Agency. Substance Infocard for butyl acrylate (CAS RN 141-32-2). Last updated 9-Sept-2023. <https://echa.europa.eu/de/substance-information/-/substanceinfo/100.004.983>
- ED Lists (2022). Endocrine Disruptor Lists. Last updated Apr-2022. <https://edlists.org/the-ed-lists>
- EFSA. European Food Safety Authority. Accessed Oct-2023. <https://www.efsa.europa.eu/en>
- EMA. European Medicines Agency. Medicines Database. Accessed Mmm-YYYY. <https://www.ema.europa.eu/en/medicines>
- EPA (a). US Environmental Protection Agency. CompTox Chemicals Dashboard. Physchem Prop. Record for butyl acrylate (CAS RN 141-32-2). Accessed Oct-2023. <https://comptox.epa.gov/dashboard/chemical/properties/DTXSID6024676>
- EPA (b). US Environmental Protection Agency. Integrated Risk Information System (IRIS). Accessed Oct-2023. https://iris.epa.gov/AtoZ/?list_type=alpha
- EU (2018). European Union. Regulation (EU) No 1169/2011 of the European Parliament and of the Council of 25 October 2011 on the provision of food information to consumers, amending Regulations (EC) No 1924/2006 and (EC) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the

Additive Toxicity Statement

2-Propenoic acid, butyl ester



Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation (EC) No 608/2004. Current consolidated version 01-Jan-2018. Document 02011R1169-20180101, Annex II. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A02011R1169-20180101>

FDA (2023). US Food & Drug Administration. Substances Added to Food inventory. Last updated 24-Aug-2023. <https://www.accessdata.fda.gov/scripts/fdcc/?set=FoodSubstances>
FEMA Flavor Ingredient Library. Flavor and Extract Manufacturers Association of the United States. Accessed Oct-2023. <https://www.femaflavor.org/flavor-library>

GESTIS (2023). Institute for Occupational Safety and Health of the German Social Accident Insurance Database. Last updated April-2023. <https://limitvalue.ifa.dguv.de/>

GSC. Good Scents Company. Record for butyl acrylate (CAS RN 141-32-2). Accessed Oct-2023. www.thegoodscentscompany.com/data/rw1226061.html

Health Canada (2018). Health Canada and Environment and Climate Change Canada. Screening Assessment. Acrylates and methacrylates group. September 2018.

<https://www.canada.ca/content/dam/eccc/documents/pdf/pded/acrylates-methacrylates/Screening-assessment-acrylates-methacrylates.pdf>

HSDB (2010). Hazardous Substances Data Bank. Record for n-butyl acrylate (CAS RN 141-32-2). Last revision date 09-Feb-2010. <https://pubchem.ncbi.nlm.nih.gov/source/hsdb/305>

IARC (2023). International Agency for Research on Cancer. List of classifications, Volumes 1–134. Last updated 11-Oct-2023. <https://monographs.iarc.who.int/list-of-classifications>

JECFA. Joint FAO/WHO Expert Committee on Food Additives. Portal. Accessed Oct-2023. <https://apps.who.int/food-additives-contaminants-jecfa-database/>

Murphy S, Ellis-Hutchings R, Finch L, Welz S and Wiench K (2018). *In vitro* genotoxicity studies: n-Butyl acrylate L5178Y mouse lymphoma (TK+/- locus assay), 2-ethylhexyl acrylate gene mutation assay in Chinese hamster V79 cells, and 2-ethylhexyl acrylate micronucleus test in human lymphocytes. Data in Brief 20, 316-325.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6111055/>

NTP (2021). National Toxicology Program. Fifteenth Report on Carcinogens (RoC). Released 21-Dec-2021. <https://ntp.niehs.nih.gov/whatwestudy/assessments/cancer/roc/index.html>

OECD (2002). Organisation for Economic Co-operation and Development. Addendum to SIDS initial assessment report: n-butyl acrylate: CAS No. 141-32-2. SIAM [SIDS Initial Assessment Meeting] 15, 22-25 October 2002. <https://hpvchemicals.oecd.org/ui/handler.axd?id=4e32e8ca-a718-4300-b753-e1a079c33900> (also cited in Health Canada, 2018).

OEHHA (2023). California Office of Environmental Health Hazard Assessment) Proposition 65 list. Dated 11-Aug-2023. <https://oehha.ca.gov/proposition-65/proposition-65-list>

PubChem (2023). An open chemistry database at the National Institutes of Health (NIH). Record for butyl acrylate (CAS RN 141-32-2). Created 16-Sept-2004. Last modified 3-Nov-2023. <https://pubchem.ncbi.nlm.nih.gov/compound/8846>

Schulz M (2016). BASF SE. n-Butyl acrylate. *In vitro* gene mutation test in L5178Y mouse lymphoma cells, OECD TG 490. BASF SE, Experimental Toxicology and Ecotoxicology, Germany “as cited in the ECHA dissemination dossier n-butyl acrylate” (cited in Murphy *et al.*, 2018).

SCOEL (1993). Recommendations of the Scientific Committee on Occupational Exposure Limits for n-butylacrylate.

https://echa.europa.eu/documents/10162/35144386/038_butyl_acrylate_oel_en.pdf

Suh M, Proctor D, Chappell G, Rager J, Thompson C, Borghoff S, Finch L, Ellis-Hutchings R and Wiench K (2018). A review of the genotoxic, mutagenic, and carcinogenic potentials of several lower acrylates. Toxicology 402-403, 50-67.

Additive Toxicity Statement

2-Propenoic acid, butyl ester



Toxtree. Estimation of toxic hazard – a decision tree approach. Ideaconsult Ltd. (Version 3.1.0). Accessed Oct-2023. Available for download at: <http://toxtree.sourceforge.net/>
WAO (2019). World Allergy Organization. Quirce S and Sastre J. Sensitizing agents inducers of occupational asthma, hypersensitivity pneumonitis and eosinophilic bronchitis. Updated Jul-2019. <http://www.worldallergy.org/education-and-programs/education/allergic-disease-resource-center/professionals/occupational-allergens/>

APPENDIX

Statement:

This ingredient has been assessed by BAT for its toxicity and carcinogenic, mutagenic or toxic for reproduction (CMR) properties. Based on an investigation of available and relevant internal and external data, it has been concluded that this ingredient does not have any CMR properties. Furthermore, this ingredient is not subject to a harmonised classification as CMR under CLP (1272/2008).