

DESCRIPTION

Name Citral

IUPAC name

CAS 5392-40-5 (2E)-3,7-dimethylocta-2,6-dienal (2Z)-3,7-dimethylocta-2,6-dienal

CAS no 5392-40-5

106-26-3

EINECS no.

CAS 5392-40-5 226-394-6 CAS 106-26-3 203-379-2

Molecular formula $C_{10}H_{16}O$

Structural formula

CAS 5392-40-5

CH, CH, H

CAS 106-26-3

H₃C CH₃

Molecular weight (g/mol.) 152.23

CLP HEALTH HAZARD CLASSIFICATION

This ingredient has been classified under Regulation (EC) No 1272/2008 of the European Parliament and of the Council and is in the classification and labelling inventory.

The CLP classifications of this ingredient are not considered applicable for one, or more, of the following reasons:

• the level required to elicit a response in a toxicological study, with a relevant route of exposure, is far higher than that of its use in BAT products



- the toxicological effect identified does not relate to the route of exposure associated with its use in BAT products
- no relevant toxicological study was identified relating to the route of exposure, and there
 is only limited consumer exposure to this ingredient associated with its use in BAT
 products

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are: TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this, our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch $et\ al^{11}$.

In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in eliquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴



Assessment of the irritation potential of Swedish snus ingredients using the Epioral[™] tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself.

In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

- 1. Available at: http://www.bibra-information.co.uk/supported access to our chemical toxicology database TRACE.html
- 2. Available at: http://toxnet.nlm.nih.gov/index.html
- 3. Available at: http://ccinfoweb.ccohs.ca/rtecs/search.html
- 4. Available at: http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384
- Available at: http://www.inchem.org/
- 6. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
- 7. Available at: http://echa.europa.eu/information-on-chemicals
- 8. Available at: http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing
- 9. Available at: http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp
- 10. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
- 11. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
- 12. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. Food Chem. Toxicol. 42 Suppl: S53-S83, 2004.
- 13. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in eliquids. Regul. Toxicol. Pharmacol. 72 (2):361-369, 2015.
- B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. Toxicology Letters 229S:S109, 2014. (Abstract)
- 15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. The Toxicologist, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
- World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans.
 Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004



- 17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific.committees/emerging/docs/scenihr_o_029.pdf.
- http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_029.pdf.
 SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_051.pdf.



REFERENCES

Alfa Aesar

Accessed February 2012. Available at: http://www.alfa.com/en/go160w.pgm?srchtyp=product

Brunn Poulsen, P. and Strandesen, M., 2011

Survey and health assessment of cosmetic products marketed as "non-preserved". Danish Ministry of the Environment. Survey of Chemical Substances in Consumer Products No. 111 2011. http://www2.mst.dk/udgiv/publications/2011/04/978-87-92708-67-0.pdf.

Burdock, G.A., 2010

Fenaroli's Handbook of Flavor Ingredients. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.

ChemIDplus

Accessed February 2012. Available at: http://chem.sis.nlm.nih.gov/chemidplus/

ChemSpider

Royal Society of Chemistry chemical structure database. Records for Citral and Citral B (5392-40-5 and 106-26-3). Accessed February 2012. Available at: http://www.chemspider.com/

CoE, 2008

Natural sources of flavourings. Report No. 3. Council of Europe Publishing. ISBN 978-92-871-6422-3.

CoE. 2007

Natural sources of flavourings. Report No. 2. Council of Europe Publishing. ISBN 978-92-871-6156-7.

CoE. 2000

Chemically-defined flavouring substances. Council of Europe Publishing. ISBN 92-871-4453-2.

CoE, 1981

Flavouring substances and natural sources of flavourings. Council of Europe. ISBN 2-7160-0081-6.

Connor, M.J., 1991

Modulation of tumor promotion in mouse skin by the food additive citral (3,7-dimethyl-2,6-octadienal). *Cancer Lett.*, 56, 25-28.

de Groot, A.C., Coenraads, P.J., Bruynzeel, D.P., Jagtman, B.A., van Ginkel, C.J., Noz, K., van der Valk, P.G., Pavel, S., Vink, J., Weyland, J.W., 2000

Routine patch testing with fragrance chemicals in the Netherlands. *Contact Dermatitis*, 42, 184-185.

De-Oliveira, A.C.A.X., Ribeiro-Pinto, L.F., Paumgartten, J.R., 1997

In vitro inhibition of CYP2B1 monooxygenase by beta-myrcene and other monoterpenoid compounds. *Toxicology Lett.*, 92, 39-46.



Duerksen-Hughes, P.J., Yang, J., Ozcan, O., 1999

p53 Induction as a genotoxic test for twenty-five chemicals undergoing *in vivo* carcinogenicity testing. *Envir. Hlth Perspect.*, 107, 805-812.

ECHA

European Chemicals Agency. Information on Chemicals. Accessed February 2012. Available at: http://echa.europa.eu/information-on-chemicals

ECHA, 2013

European Chemicals Agency. Classification and Labelling (C&L) Inventory database. Last updated 28 June 2013. Available at: http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database

ECHA, 2011

Information on registered substances. Citral. Searched 23 February 2012. http://apps.echa.europa.eu/registered/data/dossiers/DISS-9eae68e6-a212-5d5b-e044-00144f67d031/DISS-9eae68e6-a212-5d5b-e044-00144f67d031_DISS-9eae68e6-a212-5d5b-e044-00144f67d031.html.

EFSA

European Food Safety Authority. Accessed February 2012. Available at: http://www.efsa.europa.eu/

EPA

US Environmental Protection Agency. Integrated Risk Information System (IRIS). Accessed July 2013. http://www.epa.gov/iris/search_human.htm and http://cfpub.epa.gov/ncea/iris_drafts/erd.cfm?excCol=Archive&archiveStatus=both

EPISuite

Accessed February 2012. The database is available to download at: http://www.epa.gov/opptintr/exposure/pubs/episuitedl.htm

Ezendam, J., de Klerk, A., Vermeulen, J., Fokkens, P.H.B., van Loveren, H., 2009 Immune effects of inhalation exposure to fragrance allergens. RIVM Report 340301003/2009. http://www.rivm.nl/bibliotheek/rapporten/340301003.html.

Ezendam, J.W., ter Burg, W., Wijnhoven, S.W.P., 2011
Inhalation exposure to fragrance allergens. Are consumers at risk for respiratory allergies?
RIVM Report 340301004/2011. Ministry of Health, Welfare and Sport, Dutch National Institute for Public Health and the Environment (RIVM).

http://www.rivm.nl/bibliotheek/rapporten/340301004.pdf.

FDA, 2011

Food and Drug Administration. Everything added to Food in the United States (EAFUS). Last updated 17 November 2011. Available at: http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing

Fisher Scientific

Accessed February 2012. Available at: http://www.fisher.co.uk/index.php/en/technical-support



Flavis

EUROPA food flavouring website. Accessed February 2012. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm

Frosch, P.J., Rastogi, S., Pirker, C., Brinkmeier, T., Andersen, K.E., Bruze, M., Svedman, C., Goossens, A., White, I.R., Uter, W., Giménez Arnau, E., Lepoittevin, J-P., Duus Johansen, J., Menné, T., 2005

Patch testing with a new fragrance mix - reactivity to the individual constituents and chemical detection in relevant cosmetic products. *Contact Dermatitis*, 52, 216.

Frosch, P., Pirker, C., Rastogi, S., Andersen, K.E., Bruze, M., Goossens, A., White, I.R., Uter, W., Duus Johansen, J., Menné, T., 2004

The new fragrance mix II - test results of a multicentre European study. *Contact Dermatitis*, 50, 149

Frosch, P.J., Johansen, J.D., Menné, T., Pirker, C., Rastogi, S.C., Andersen, K.E., Bruze, M., Goossens, A., Lepoittevin, J.P., White, I.R., 2002

Further important sensitizers in patients sensitive to fragrances. I. Reactivity to 14 frequently used chemicals. *Contact Dermatitis*, 47, 78-85.

Frosch, P.J., Pilz, B., Andersen, K.E., Burrows, D., Camarasa, J.G., Dooms-Goossens, A., Ducombs, G., Fuchs, T., Hannuksela, M., Lachapelle, J.M., Lahti, A., Maibach, H.I., Menné, T., Rycroft, R.J.G., Shaw, S., Wahlberg, J.E., White, I.R., Wilkinson, J.D., 1995

Patch testing with fragrances: results of a multicenter study of the European Environmental and Contact Dermatitis Research Group with 48 frequently used constituents of perfumes. *Contact Dermatitis*, 33, 333-342.

GESTIS

Gefahrstoffinformationssystem (Databases on hazardous substances). Accessed February 2012. Available at: http://limitvalue.ifa.dguv.de/Webform_gw.aspx

Golomb, E., Scolnik, M., Koren, R., Servadio, C., Sandbank, U., Abramovici, A., 2001 Effects of senescence and citral on neuronal vacuolar degeneration in rat pelvic ganglia. *Neurotoxicology*, 22, 73-77.

Gomes-Carneiro, M.R., Felzenszwalb, I., Paumgartten, F.J., 1998

Mutagenicity testing (±)-camphor, 1,8-cineole, citral, citronellal, (-)-menthol and terpineol with the Salmonella/microsome assay. *Mutation Res.*, 416, 129-136.

Good Scents Company

Accessed February 2012. Available at: http://www.thegoodscentscompany.com/

Heisterberg, M.V., Menné, T., Johansen, J.D., 2011

Contact allergy to the 26 specific fragrance ingredients to be declared on cosmetic products in accordance with the EU cosmetics directive. *Contact Dermatitis*, 65, 266-275.

Heydorn, S., Duus Johansen, J., Andersen, K.E., Basketter, D.A., Bruze, M., Karlberg, A.T., White, I.R., Menné, T., 2002

Identification of fragrances relevant to hand eczema. Contact Dermatitis, 46(Suppl.4), 21-22.



Heydorn, S., Duus Johansen, J., Andersen, K.E., Bruze, M., Svedman, C., White, I.R., Basketter, D.A., Menné, T., 2003a

Fragrance allergy in patients with hand eczema – a clinical study. *Contact Dermatitis*, 48, 317-323.

Heydorn, S., Menné, T., Andersen, K.E., Bruze, M., Svedman, C., White, I.R., Basketter, D.A., 2003b

Citral a fragrance allergen and irritant. Contact Dermatitis, 49, 32-36.

Hoberman, A.M., Christian, M.S., Bennett, M.M., Vollmuth, T.A., 1989 Oral general reproduction study of citral in female rats. *Toxicologist*, 9, 271.

HSDB. 2007

Hazardous Substances Databank. Record for Citral, number: 993. Last revision date: 11 October 2007. http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB.

IARC

International Agency for Research on Cancer. World Health Organization. Accessed July 2013. Available at: http://www.iarc.fr/

Ishidate, M., Sofuni, T., Yoshikawa, K., Hayashi, M., Nohmi, T., Sawada, M., Matsuoka, A., 1984

Primary mutagenicity screening of food additives currently used in Japan. *Fd Chem. Toxic.*, 22, 623-636.

Itoh, M., Ishihara, M., Hosono, K., Kantoh, H., Kinoshita, M., Yamada, K., Nishimura, M., 1986 Results of patch tests conducted between 1978 and 1985 using cosmetic ingredients. *Skin Res.*, 28, 110-119.

JECFA

Joint Expert Committee on Food Additives. Monographs and Evaluations. Accessed February 2012. Available at: http://www.inchem.org/pages/jecfa.html

JECFA, 2004a

Evaluation of certain food additives and contaminants. Sixty-first report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series, no. 922. World Health Organization. Geneva.

http://whqlibdoc.who.int/trs/WHO_TRS_922.pdf.

JECFA, 2004b

Safety evaluation of certain food additives and contaminants. Aliphatic branched-chain saturated and unsaturated alcohols, aldehydes, acids, and related esters. Prepared by the Sixty-first meeting of the Joint FAO/WHO Expert Committee on Food Additives. WHO Food Additives Series, No. 52. World Health Organization, Geneva. http://whqlibdoc.who.int/publications/2004/924166052X.pdf.



JECFA, 1999

Evaluation of certain food additives and contaminants. Forty-ninth report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series, No. 884. http://whqlibdoc.who.int/trs/WHO_TRS_884.pdf

JECFA. 1980

Evaluation of certain food additives. Twenty-third report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series, No. 648, and corrigenda. http://whqlibdoc.who.int/trs/WHO_TRS_648.pdf

JRC

Joint Research Centre. European chemical Substances Information System (ESIS). Accessed February 2012. Available at: http://esis.jrc.ec.europa.eu/

Lalko, J. and Api, A.M., 2008

Citral: identifying a threshold for induction of dermal sensitization. *Regul. Toxic. Pharmac.*, 52, 62-73.

Lalko, J. and Api, A.M., 2006

Investigation of the dermal sensitization potential of various essential oils in the local lymph node assay. *Fd Chem. Toxic.*, 44, 739-46.

Merck, 2006

The Merck Index. An Encyclopedia of Chemicals, Drugs and Biologicals. Ed. O'Neil J et al. Fourteenth edition. Merck & Co., Ltd., Whitehouse Station, NJ, USA.

Nogueira, A.C.M.A., Carvalho, R.R., Souza, C.A., Chahoud, I., Paumgartten, F.J., 1995 Study on the embryofeto-toxicity of citral in the rat. *Toxicology*, 96, 105-113.

NTP

US National Toxicology Program. Accessed July 2013. Available at http://ntp.niehs.nih.gov/?objectid=03C9AF75-E1BF-FF40-DBA9EC0928DF8B15 and http://tools.niehs.nih.gov/ntp_tox/index.cfm

NTP, 2003

NTP technical report on the toxicology and carcinogenesis studies of citral (microencapsulated) (CAS no. 5392-40-5) in F344/n rats and B6C3F1 mice (feed studies). NTP TR 505 NIH Publication No. 03-4439. National Toxicology Program January 2003. http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr505.pdf.

OECD, 2001

SIDS Initial Assessment Report for 13th SIAM. Citral. Switzerland, November 6-9, 2001. http://www.chem.unep.ch/irptc/sids/oecdsids/5392-40-5.pdf.

PubChem

Accessed February 2012. Available at: http://pubchem.ncbi.nlm.nih.gov/



Ress, N.B., Hailey, J.R., Maronpot, R.R., Bucher, J.R., Travlos, G.S., Haseman, J.K., Orzech, D.P., Johnson, J.D., Hejtmancik, M.R., 2003

Toxicology and carcinogenesis studies of microencapsulated citral in rats and mice. *Toxic. Sci.*, 71, 198-206.

SCENIHR, 2010

Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific committees/emerging/docs/scenihr o 029.pdf

Schnuch, A., Uter, W., Geier, J., Lessmann, H., Frosch, P.J., 2007 Sensitization to 26 fragrances to be labelled according to current European regulation Results of the IVDK and review of the literature. *Contact Dermatitis*, 57, 1-10.

Sigma-Aldrich

Accessed February 2012. Available at: http://www.sigmaaldrich.com/united-kingdom.html

SRC, 2011

Syracuse Research Corporation. Interactive PhysProp Database Demo. Record for Citral. Available at: http://www.syrres.com/what-we-do/databaseforms.aspx?id=386

Steltenkamp, R.J., Booman, K.A., Dorsky, J., King, T.O., Rothenstein, A.S., Schwoeppe, E.A., Sedlak, R.I., Smith, T.H., Thompson, G.R., 1980

Citral: A survey of consumer patch-test sensitization. Fd Cosmet. Toxicol., 18, 413-418.

Tapsell, L.C., Hemphill, I., Cobiac, L., Patch, C.S., Sullivan, D.R., Fenech, M., Roodenrys, S., Keogh, J.B., Clifton, P.M., Williams, P.G., Fazio, V.A., Inge, K.E., 2006

Health benefits of herbs and spices: the past, the present, the future. *Med. J. Aust.*, 185(Suppl. 4), S1-S24.

http://www.mja.com.au/public/issues/185_04_210806/supplement_210806.html

Toaff, M.E., Abramovici, A., Sporn, J., Liban, E., 1979

Selective oocyte degeneration and impaired fertility in rats treated with aliphatic monoterpene, Citral. *J. Reprod. Fert.*, 55, 347-352.

York, R.G., Vollmuth, T.A., Gaworski, C.L., 1989

Developmental toxicity evaluation of inhaled citral in rats. *Toxicologist*, 9, 271.

Yoshimura, H., Shigeno, H., Kawamura, K., Furukawa, M., undated

Preliminary reproduction toxicity screening test of citral by oral administration in rats. Safety Research Institute for Chemical Compounds Co., Ltd. 363-24 Shin-ei, Kiyota-ku, Sapporo, Hokkaido, 004-0839, Japan.

http://dra4.nihs.go.jp/mhlw_data/jsp/ResultPageENG.jsp?condition_item=CAS%94%D4%8D%8 6&condition_keyword=5392-40-5&condition_type=*.

Zeiger, E., Anderson, B., Haworth, S., Lawlor, T., Mortelmans, K., Speck, W., 1987 Salmonella mutagenicity tests: III. Results from the testing of 255 chemicals. *Envir. molec. Mutagen.*, 9 (Suppl.9), 1-110.