


Additive Toxicity Statement

Ethyl vanillin

DESCRIPTION

Name	Ethyl vanillin
IUPAC name	3-Ethoxy-4-hydroxybenzaldehyde.
CAS no	121-32-4
EINECS no.	204-464-7
Molecular formula	C ₉ H ₁₀ O ₃
Structural formula	



Molecular weight (g/mol.)	166.17
---------------------------	--------

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are: TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Additive Toxicity Statement

Ethyl vanillin



Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this, our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch *et al*¹¹.

In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴
- Assessment of the irritation potential of Swedish snus ingredients using the Epioral™ tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself.

In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

1. Available at: <http://www.bibra-information.co.uk/supported> access to our chemical toxicology database TRACE.html
2. Available at: <http://toxnet.nlm.nih.gov/index.html>
3. Available at: <http://ccinfoweb.ccohs.ca/rtecs/search.html>
4. Available at: <http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384>
5. Available at: <http://www.inchem.org/>
6. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
7. Available at: <http://echa.europa.eu/information-on-chemicals>
8. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>
9. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp>
10. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
11. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
12. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem. Toxicol.* 42 Suppl: S53-S83, 2004.
13. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids. *Regul. Toxicol. Pharmacol.* 72 (2):361-369, 2015.
14. B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. *Toxicology Letters* 229S:S109, 2014. (Abstract)
15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. *The Toxicologist*, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
16. World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004
17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf.
18. SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_051.pdf.

REFERENCES

Adams, T.B., Cohen, S.M., Doull, J., Feron, V.J., Goodman, J.I., Marnett, L.J., Munro, I.C., Portoghese, P.S., Smith, R.L., Waddell, W.J., Wagner, B.M; Expert Panel of the Flavor and Extract Manufacturers Association, 2005

The FEMA GRAS assessment of hydroxyl- and alkoxy-substituted benzyl derivatives used as flavor ingredients. *Fd Chem. Toxic.*, 43, 1241-1271.

Basketter, D.A., Wright, Z.M., Warbrick, E.V., Dearman, R.J., Kimber, I., Ryan, C.A., Gerberick, G.F., White, I.R., 2001

Human potency predictions for aldehydes using the local lymph node assay. *Contact Dermatitis*, 45, 89-94.

Basketter, D.A., Evans, P., Gerberick, G.F., Kimber, I.A., 2002

Factors affecting thresholds in allergic contact dermatitis: safety and regulatory considerations. *Contact Dermatitis*, 47, 1-6.

Burdock, G.A., 2010

Fenaroli's Handbook of Flavor Ingredients. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.

CCRIS, 2006

Chemical Carcinogenesis Research Information System. Record for ethyl vanillin. Last revision date 05 April 2006. Available at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?CCRIS>

ChemIDplus

Accessed February 2013. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>

ChemSpider

Royal Society of Chemistry chemical structure database. Record for ethyl vanillin (121-32-4). Accessed February 2013. Available at: <http://www.chemspider.com/>

CoE, 2000

Chemically-defined flavouring substances. Council of Europe Publishing. ISBN 92-871-4453-2.

Deichmann, W. and Kitzmiller, K.V., 1940

On the toxicity of vanillin and ethyl vanillin for rabbits and rats. *J. Am. pharm. Ass.*, 29, 425-428.

ECHA

European Chemicals Agency. Information on chemicals. Accessed on 15 November 2013. Available at: <http://echa.europa.eu/information-on-chemicals>.

EFSA, 2012

Additive Toxicity Statement

Ethyl vanillin



European Food Safety Authority. EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids (CEF); Scientific opinion on Flavouring Group Evaluation 20, Revision 4 (FGE.20Rev4): Benzyl alcohols, benzaldehydes, a related acetal, benzoic acids, and related esters from chemical groups 23 and 30. Question No EFSA-Q-2012-00621, EFSA-Q-2012-00620, EFSA-Q-2012-00619, EFSA-Q-2012-00618, adopted on 20 November 2012. The EFSA Journal 10 (12), 2994. Available at: <http://www.efsa.europa.eu/en/efsajournal/doc/2994.pdf>

EFSA, 2011

European Food Safety Authority. EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids (CEF); Scientific opinion on Flavouring Group Evaluation 20, Revision 3 (FGE.20Rev3): Benzyl alcohols, benzaldehydes, a related acetal, benzoic acids, and related esters from chemical groups 23 and 30. Question No EFSA-Q-2011-00299, adopted on 19 May 2011. The EFSA Journal 9(7), 2176. Available at: <http://www.efsa.europa.eu/en/efsajournal/doc/2176.pdf>

EFSA, 2008

European Food Safety Authority. Flavouring Group Evaluation 52 (FGE.52): Consideration of hydroxy- and alkoxy-substituted benzyl derivatives evaluated by JECFA (57th meeting) structurally related to benzyl alcohols, benzaldehydes, a related acetal, benzoic acids, and related esters evaluated by EFSA in FGE.20 (2005). (Commission Regulation (EC) No 1565/2000 of 18 July 2000). Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in contact with Food (AFC). Question No. EFSA-Q-2008-032C. Adopted on 03 July 2007. The EFSA Journal (2008) 637, 1. Available at: http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178692189524.htm.

EPA

US Environmental Protection Agency. Integrated Risk Information System. Accessed June, 2013. Available at: http://www.epa.gov/iris/search_human.htm

EPISuite

Accessed February 2013. The database is available to download at: <http://www.epa.gov/opptintr/exposure/pubs/episuitedi.htm>

FDA, 2012

Food and Drug Administration. CFR – Code of Federal Regulations Title 21. Last updated 1 April 2012. Available at: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/cfrsearch.cfm>

FDA, 2011

Food and Drug Administration. Everything added to Food in the United States (EAFUS). Last updated 17 November 2011. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>

Flavis

Flavouring substances data-base. Accessed on 22 November 2013. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm?fullTextSearch=&flavisname=Benzoin&eugroup=all&Evaluation=all&Evaluationnumber=&inreg=-1&FLNO=&CASNO=&JECFANO=&FEMANO=&CENO.

Additive Toxicity Statement

Ethyl vanillin



Frosch, P.J., Pilz, B., Andersen, K.E., Burrows, D., Camarasa, J.G., Dooms-Goossens, A., Ducombs, G., Fuchs, T., Hannuksela, M., Lachapelle, J.M., Lahti, A., Maibach, H.I., Menné, T., Rycroft, R.J.G., Shaw, S., Wahlberg, J.E., White, I.R., Wilkinson, J.D., 1995

Patch testing with fragrances: results of a multicentre study of the European Environmental and Contact Dermatitis Research Group with 48 frequently used constituents of perfumes. *Contact Dermatitis*, 33, 333-342.

Fuglsang, G., Madsen, G., Halken, S., Jørgensen, S., Ostergaard, P.A., Osterballe, O., 1994
Adverse reactions to food additives in children with atopic symptoms. *Allergy*, 49, 31-37.

IARC

International Agency for Research on Cancer. Agents classified by the IARC Monographs, Volumes 1-108. Accessed 24 October, 2013. Available at: <http://monographs.iarc.fr/ENG/Classification/ClassificationsCASOrder.pdf>

Ishidate, M., Sofuni, T., Yoshikawa, K., Hayashi, M., Nohmi, T., Sawada, M., Matsuoka, A., 1984

Primary mutagenicity screening of food additives currently used in Japan. *Fd Chem. Toxic.*, 22, 623-636.

JECFA, 2002

Safety evaluation of certain food additives and contaminants. Hydroxy - and alkoxy-substituted benzyl derivatives. Prepared by the fifty-seventh meeting of the Joint FAO/WHO Expert Committee on Food Additives. WHO Food Additives Series 48. WHO, Geneva. Available at: <http://www.inchem.org/documents/jecfa/jecmono/v48je15.htm>

JECFA, 1996

Toxicological evaluation of certain food additives and contaminants in food. Prepared by the forty-fourth meeting of the Joint FAO/WHO Expert Committee on Food Additives. WHO Food Additives Series 35. WHO, Geneva. Available at: <http://www.inchem.org/documents/jecfa/jecmono/v35je07.htm>

Jenner, P.M., Hagan, E.C., Taylor, J.M., Cook, E.L., Fitzhugh, O.G., 1964

Food flavourings and compounds of related structure. I. Acute oral toxicity. *Food Cosmet. Toxicol.*, 2, 327-343.

JRC

Joint Research Centre. European chemical Substances Information System (ESIS). Accessed February 2013. Available at: <http://esis.jrc.ec.europa.eu/>

Merck, 2006

The Merck Index. An Encyclopedia of Chemicals, Drugs and Biologicals. Ed. O'Neil J et al, Fourteenth edition. Merck & Co., Ltd., Whitehouse Station, NJ, USA.

Miller, D., Wheals, B.B., Beresford, N., Sumpter, J.P., 2001

Estrogenic activity of phenolic additives determined by an *in vitro* yeast bioassay. *Envir. Hlth Perspect.*, 109, 133.

Monsanto, 1991

Additive Toxicity Statement

Ethyl vanillin



Final report of several test with ethavan with cover letter dated 112191. Contractor Younger Laboratories Inc. Study Date 17 August 1976. EPA/OTS; Doc #86-920000149. Microfiche No. OTS0534355.

Mortelmans, K., Haworth, S., Lawlor, T., Speck, W., Tainer, B., Zeiger, E., 1986
Salmonella mutagenicity tests: II. Results from the testing of 270 chemicals. *Envir. Mutagen.*, 8. Suppl. 7, 1-119.

NTP, 2011

National Toxicology Programme. Twelfth report on carcinogens. Accessed June, 2013. Available at: <http://ntp.niehs.nih.gov/?objectid=03C9AF75-E1BF-FF40-DBA9EC0928DF8B15>.

NTP, 1996

National Toxicology Program: Annual plan for fiscal year 1996. PB97141675.

NTP, 1994

National Toxicology Program. Study A38110 (Micronucleus). Available at: http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm

NTP, 1982

National Toxicology Program. Study 340491 (Salmonella). Available at: http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm

Oda, Y., Hamono, Y., Inoue, K., Yamamoto, H., Niihara, T., Kunita, N., 1978
Mutagenicity of food flavours in bacteria. 1. *Shouhin Eisei Hen*, 9, 177-181.

Ohuchida, A., Furukawa, A., Yoshida, R., 1989

Micronucleus test of polyploidy inducers. *Mutation Res.*, 216, 371-372.

Opdyke, D.L.J., 1975

Monographs on Fragrance Raw Materials: Ethyl vanillin. *Fd Cosmet. Toxicol.*, 13, 103-104.

Patlewicz, G., Basketter, D.A., Smith, C.K., Hotchkiss, S.A., Roberts, D.W., 2001

Skin-sensitization structure-activity relationships for aldehydes. *Contact Dermatitis*, 44, 331-336.

Rudzki, E. and Grzywa, Z., 1976

Immediate reactions to balsom of Peru, cassia oil and ethyl vanillin. *Contact Dermatitis*, 2, 360-361.

Sasaki, Y.F., Imanishi, H., Ohta, T., Shirasu, Y., 1987

Effects of antimutagenic flavourings on SCEs induced by chemical mutagens in cultured Chinese hamster cells. *Mutation Res.*, 189, 313.

Additive Toxicity Statement

Ethyl vanillin



SCCS, 2012

Scientific Committee on Consumer Safety (SCCS). Opinion on fragrance allergens in cosmetic products. Opinion adopted by the SCCS at its 15th plenary meeting of 26-27 June 2012. SCCS/1459/11. Available at:

http://ec.europa.eu/health/scientific_committees/consumer_safety/docs/sccs_o_102.pdf

SCENIHR, 2010

Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at:

http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf

Sigma-Aldrich, 2013

MSDS for ethyl vanillin. Product No. W246409. Revision date: 02 March 2013. Accessed March 2013.

Available at: http://www.sigmaaldrich.com/Area_of_Interest/Europe_Home/UK.html.

Stoner, G.D., Shimkin, M.B., Kniazeff, A.J., Weisburger, J.H., Weisburger, E.K., Gori, G.B., 1973

Test for carcinogenicity of food additives and chemotherapeutic agents by the pulmonary tumor response in strain A mice. *Cancer Res.*, 33, 3069-3085.

Wild, D., King, M.T., Gocke, E., Eckhardt, K., 1983

Study of artificial flavouring substances for mutagenicity in the Salmonella/microsome, BASC and micronucleus tests. *Fd Chem. Toxic.*, 21, 707-719.