

Additive Toxicity Statement

Lemon oil, expressed



DESCRIPTION

Name Lemon oil, expressed

IUPAC name Not applicable

CAS no 8020-19-7
8008-56-8

EINECS no.
CAS 8020-19-7; 8008-56-8 Not specified.

Molecular formula
CAS 8020-19-7; 8008-56-8 Not applicable.

Structural formula
CAS 8020-19-7; 8008-56-8 Not applicable.

Molecular weight (g/mol.)
CAS 8020-19-7; 8008-56-8 Not applicable.

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are: TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this,

Additive Toxicity Statement

Lemon oil, expressed



our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch *et al*¹¹.

In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴
- Assessment of the irritation potential of Swedish snus ingredients using the Epioral™ tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself.

In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

1. Available at: [http://www.bibra-information.co.uk/supported access to our chemical toxicology database TRACE.html](http://www.bibra-information.co.uk/supported%20access%20to%20our%20chemical%20toxicology%20database%20TRACE.html)

Additive Toxicity Statement

Lemon oil, expressed



2. Available at: <http://toxnet.nlm.nih.gov/index.html>
3. Available at: <http://ccinfoweb.ccohs.ca/rtecs/search.html>
4. Available at: <http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384>
5. Available at: <http://www.inchem.org/>
6. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
7. Available at: <http://echa.europa.eu/information-on-chemicals>
8. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>
9. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp>
10. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
11. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
12. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem. Toxicol.* 42 Suppl: S53-S83, 2004.
13. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids. *Regul. Toxicol. Pharmacol.* 72 (2):361-369, 2015.
14. B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. *Toxicology Letters* 229S:S109, 2014. (Abstract)
15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. The Toxicologist, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
16. World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004
17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf.
18. SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_051.pdf.

REFERENCES

- Abifadel, R., Mortureux, P., Perromat, M., Ducombs, G., Taieb, A., 1992
Contact sensitivity to flavourings and perfumes in atopic dermatitis. *Contact Dermatitis*, 27, 43-46.
- Audicana, M. and Bernaola, G., 1994
Occupational contact dermatitis from citrus fruits: lemon essential oils. *Contact Dermatitis*, 31, 183-184.
- Brites, M.M., Gonçalo, M., Figueiredo, A., 2000
Contact allergy to fragrance mix – a 10-year study. *Contact Dermatitis*, 43, 181-182.
- Burdock, G.A., 2010
Fenaroli's Handbook of Flavor Ingredients. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.
- Ceccarelli, I., Lariviere, W.R., Fiorenzani, P., Sacerdote, P., Aloisi, A.M., 2004
Effects of long-term exposure of lemon essential oil odor on behavioral, hormonal and neuronal parameters in male and female rats. *Brain Res.*, 1001, 78-86.
- ChemIDplus
Accessed on 13 August 2012. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>
- CoE, 2000
Natural sources of flavourings. Report No. 1. Council of Europe Publishing. ISBN 92-871-4324-2.
- Crebelli, R., Aquilina, G., Conti, L., Carere, A., 1990
Microbial mutagenicity screening of natural flavouring substances. *Microbiologica*, 13, 115-119.
- ECHA, 2014
European Chemicals Agency. Classification and Labelling (C&L) Inventory database. Last updated 3 October 2014. Available at: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database>
- ECHA
European Chemicals Agency. Information on Chemicals. Accessed November 2014. Available at: <http://echa.europa.eu/information-on-chemicals>
- EFSA
European Food Safety Authority. Accessed November 2014. Available at: <http://www.efsa.europa.eu/>
- EPA
US Environmental Protection Agency. Integrated Risk Information System (IRIS). Accessed November 2014. Available at: http://www.epa.gov/iris/search_human.htm and http://cfpub.epa.gov/ncea/iris_drafts/erd.cfm?excCol=Archive&archiveStatus=both

Additive Toxicity Statement

Lemon oil, expressed



FDA, 2011

Food and Drug Administration. Everything added to Food in the United States (EAFUS). Last updated 17 November 2011. Accessed xxx 2012. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>

EPISuite

Accessed November 2014. The database is available to download at: <http://www.epa.gov/opptintr/exposure/pubs/episuitedi.htm>

Flavis

EUROPA food flavouring website. Accessed November 2014. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm

Francalanci, S., Sertoli, A., Giorgini, S., Pigatto, P., Santucci, B., Valsecchi, R., 2000
Multicentre study of allergic contact cheilitis from toothpastes. *Contact Dermatitis*, 43, 216-222.

GESTIS

Gefahrstoffinformationssystem (Databases on hazardous substances). Accessed November 2014. Available at: http://limitvalue.ifa.dguv.de/Webform_gw.aspx

Good Scents Company

Records for 8008-56-8. Accessed October 2012. Available at: <http://www.thegoodscentscompany.com/>

Guin, J.D., 2005

Use of consumer product ingredients for patch testing. *Dermatitis*, 16, 71-77.

HSDB

Hazardous Substances Data Bank. Accessed November 2014. Available at: <http://toxnet.nlm.nih.gov/newtoxnet/hsdb.htm>

IARC

International Agency for Research on Cancer. World Health Organization. Accessed November 2014. Available at: <http://www.iarc.fr/>

Ishidate, M., Sofuni, T., Yoshikawa, K., Hayashi, M., Nohmi, T., Sawada, M., Matsuoka, A., 1984

Primary mutagenicity screening of food additives currently used in Japan. *Fd Chem. Toxic.*, 22, 623-636.

JECFA

Joint Expert Committee on Food Additives. Monographs and Evaluations. Accessed November 2014. Available at: <http://www.inchem.org/pages/jecfa.html>

Kuroda, K., Yoo, Y.S., Ishibashi, T., 1989

Rec-assay on natural food additives. *Seikatsu Eisei*, 33, 15-23.

Lam, L.K.T. and Zheng, B., 1991

Additive Toxicity Statement

Lemon oil, expressed



Effects of essential oils on glutathione S-transferase activity in mice. *J. agric. Fd Chem.*, 39, 660-662.

Merck, 2013

The Merck Index. An Encyclopedia of Chemicals, Drugs and Biologicals. Ed. O'Neil MJ et al, Fifteenth edition. Royal Society of Chemistry, Cambridge, UK.

Naganuma, M., Hirose, S., Nakayama, Y., Nakajima, K., Someya, T., 1985

A study of the phototoxicity of lemon oil. *Archs dermat. Res.*, 278, 31-36.

NTP

US National Toxicology Program. Accessed November 2014. Available at: <http://www.niehs.nih.gov/research/resources/databases/cebs/index.cfm> and <http://tools.niehs.nih.gov/cebs3/ui/>

NTP, 2002

US National Toxicology Program. Salmonella study on Lemon oil, CASRN 8008-56-8, study ID A40902. Available at: <http://ntp.niehs.nih.gov/>.

Opdyke, D.L.J., 1974a

Monographs on fragrance raw materials: Lemon oil expressed. *Fd Cosmet. Toxicol.*, 12, 725-726.

Opdyke, D.L.J., 1974b

Monographs on fragrance raw materials: Lemon oil distilled. *Fd Cosmet. Toxicol.*, 12, 727.

PubChem

US National Center for Biotechnology Information. Accessed November 2014. Available at: <http://pubchem.ncbi.nlm.nih.gov/>

Roe, F.J.C. and Pierce, W.E.H., 1960

Tumor promotion by citrus oils: tumors of the skin and urethral orifice in mice. *J. natn. Cancer Inst.*, 24, 1389-1403.

RTECS, 2008

Registry of Toxic Effects of Chemical Substances. Record for Lemon oil, CAS No. 8008-56-8. Last updated November 2008.

SCENIHR, 2010

Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf

Schwartz, L., 1938

Cutaneous hazards in the citrus fruit industry. *Archs Derm.*, 37, 631-649.

Serrano, G., Pujol, C., Cuadra, J., Gallo, S., Aliaga, A., 1989

Riehl's melanosis: Pigmented contact dermatitis caused by fragrances. *J. Am. Acad. Derm.*, 21, 1057-1060

Additive Toxicity Statement

Lemon oil, expressed



Sharp, D.W., 1978

The sensitization potential of some perfume ingredients tested using a modified Draize procedure. *Toxicology*, 9, 261-271.

Sigma-Aldrich, 2011

Material Safety Data Sheet for Lemon oil (CAS-No. 8008-56-8). Version 4.1 Revision date 6 December 2011. Available at: <http://www.sigmaaldrich.com/united-kingdom.html>

SRC, 2013

Syracuse Research Corporation. Interactive PhysProp Database Demo. Available at: <http://esc.syrres.com/fatepointer/search.asp>

Thomson, K.F. and Wilkinson, S.M., 2000

Allergic contact dermatitis to plant extracts in patients with cosmetic dermatitis. *Br. J. Derm.*, 142, 84-88.

Toxtree

Estimation of toxic hazard – a decision tree approach. Ideaconsult Ltd. (Version 2.6.6). Accessed November 2014. Available for download at: http://ihcp.jrc.ec.europa.eu/our_labs/computational_toxicology/qsar_tools/toxtree

Trattner, A., David, M., Lazarov, A., 2008

Occupational contact dermatitis due to essential oils. *Contact Dermatitis*, 58, 282-284.

Uter, W., Schmidt, E., Geier, J., Lessmann, H., Schnuch, A., Frosch, P., 2010

Contact allergy to essential oils: current patch test results (2000-2008) from the Information Network of Departments of Dermatology (IVDK). *Contact Dermatitis*, 63, 277-283.

Wattenberg, L.W. and Coccia, J.B., 1991

Inhibition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone carcinogenesis in mice by d-limonene and citrus fruit oils. *Carcinogenesis*, 12, 115-117.

Wöhr, S., Hemmer, W., Focke, M., Götz, M., Jarisch, R., 2001

The significance of fragrance mix, balsam of Peru, colophony and propolis as screening tools, in the detection of fragrance allergy. *Br. J. Derm.*, 145, 268-273.