

Additive Toxicity Statement

Methyl cyclopentenolone




DESCRIPTION

Name	Methyl cyclopentenolone
IUPAC name	
CAS 80-71-7	2-Hydroxy-3-methylcyclopent-2-en-1-one
CAS 765-70-8	3-Methylcyclopentane-1,2-dione
CAS no	80-71-7 (enol) 765-70-8 (keto)


EINECS no.	
CAS 80-71-7	201-303-2
CAS 765-70-8	212-154-8

Molecular formula	C ₆ H ₈ O ₂
-------------------	--

Structural formula
CAS 80-71-7



CAS 765-70-8



Molecular weight (g/mol.)	112.127
---------------------------	---------

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are: TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this, our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch *et al*¹¹.

In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴
- Assessment of the irritation potential of Swedish snus ingredients using the Epioral™ tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself.

In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

1. Available at: [http://www.bibra-information.co.uk/supported access to our chemical toxicology database TRACE.html](http://www.bibra-information.co.uk/supported%20access%20to%20our%20chemical%20toxicology%20database%20TRACE.html)
2. Available at: <http://toxnet.nlm.nih.gov/index.html>
3. Available at: <http://ccinfoweb.ccohs.ca/rtecs/search.html>
4. Available at: <http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384>
5. Available at: <http://www.inchem.org/>
6. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
7. Available at: <http://echa.europa.eu/information-on-chemicals>
8. Available at: <http://www.accessdata.fda.gov/scripts/cfn/fcnNavigation.cfm?rpt=eafusListing>
9. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp>
10. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
11. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
12. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem. Toxicol.* 42 Suppl:S53-S83, 2004.
13. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids. *Regul. Toxicol. Pharmacol.* 72 (2):361-369, 2015.
14. B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. *Toxicology Letters* 229S:S109, 2014. (Abstract)
15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. *The Toxicologist*, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
16. World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004
17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_029.pdf.
18. SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_051.pdf.

Additive Toxicity Statement

Methyl cyclopentenolone



REFERENCES

Aeschbacher, H.U., Wolleb, U., Löliger, J., Spadone, J.C., Liardon, R., 1989
Contribution of coffee aroma constituents to the mutagenicity of coffee. *Fd Chem. Toxic.*, 27, 227-232.

Alfa Aesar

Accessed March 2013. Available at: <http://www.alfa.com/en/go160w.pgm?srchtyp=product>

Bjeldanes, L.F. and Chew, H., 1979

Mutagenicity of 1,2-dicarbonyl compounds: maltol, kojic acid, diacetyl and related substances. *Mutation Res.*, 67, 367-371.

Burdock, G.A., 2010

Fenaroli's Handbook of Flavor Ingredients. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.

ChemIDplus

Accessed: March 2013. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>

ChemSpider

Royal Society of Chemistry chemical structure database. Accessed March 2013. Available at: <http://www.chemspider.com/>

CoE, 2000

Chemically-defined flavouring substances. Council of Europe Publishing. ISBN 92-871-4453-2.

EFSA, 2014

European Food Safety Authority. Scientific Opinion on Flavouring Group Evaluation 213, Revision 1 (FGE.213Rev1): Consideration of genotoxic potential for α,β -Unsaturated Alicyclic ketones and precursors from chemical subgroup 2.7 of FGE.19. EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids. *EFSA J.*, 12(5), 3661. Available at <http://www.efsa.europa.eu/en/search/doc/3661.pdf>

EFSA

European Food Safety Authority. Accessed December 2014. Available at: <http://www.efsa.europa.eu/>

EPA

US Environmental Protection Agency. Integrated Risk Information System (IRIS). Accessed December 2014. Available at: http://www.epa.gov/iris/search_human.htm and http://cfpub.epa.gov/ncea/iris_drafts/erd.cfm?excCol=Archive&archiveStatus=both

EPISuite

Accessed March 2013. The database is available to download at: <http://www.epa.gov/opptintr/exposure/pubs/episuitedi.htm>

FDA, 2011

Additive Toxicity Statement

Methyl cyclopentenolone



Food and Drug Administration. Everything added to Food in the United States (EAFUS). Last updated 17 November 2011. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>

Fisher Scientific

Accessed March 2013. Available at: <http://www.fisher.co.uk/index.php/en/technical-support>

Flavis

EUROPA food flavouring website. Accessed March 2013. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm

GESTIS

Gefahrstoffinformationssystem (Databases on hazardous substances). Accessed March 2013. Available at: http://limitvalue.ifa.dguv.de/Webform_gw.aspx

Good Scents Company

Records for cyclotene (765-70-8) and cyclotene hydrate (80-71-7). Accessed March 2013. Available at: <http://www.thegoodscentscompany.com/>

Heck, J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B., Curren, R.D., 1989

An evaluation of food flavoring ingredients in a genetic toxicity screening battery. *Toxicologist*, 9, 257.

HSDB

Hazardous Substances Data Bank. Accessed March 2013. Available at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>

IARC

International Agency for Research on Cancer. World Health Organization. Accessed March 2013. Available at: <http://www.iarc.fr/>

JECFA

Joint Expert Committee on Food Additives. Monographs and Evaluations. Accessed December 2014. Available at: <http://www.inchem.org/pages/jecfa.html>

JECFA, 1999

Safety evaluation of certain food additives. Aliphatic acyclic and alicyclic alpha-diketones and related alpha-hydroxyketones. WHO Food Additive Series 42. Prepared by the fifty-first meeting of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization, Geneva. Available at: <http://www.inchem.org/documents/jecfa/jecmono/v042je20.htm>.

JRC

Joint Research Centre. European chemical Substances Information System (ESIS). Accessed March 2013. Available at: <http://esis.jrc.ec.europa.eu/>

Merck, 2006

The Merck Index. An Encyclopedia of Chemicals, Drugs and Biologicals. Ed. O'Neil J et al, Fourteenth edition. Merck & Co., Ltd., Whitehouse Station, NJ, USA.

Additive Toxicity Statement

Methyl cyclopentenolone



NTP

US National Toxicology Program. Accessed December 2014. Available at: <http://www.niehs.nih.gov/research/resources/databases/cebs/index.cfm> and <http://tools.niehs.nih.gov/cebs3/ui/>

Opdyke, D.L.J., 1976

Monographs on Fragrance Raw Materials: Methylcyclopentenolone. *Fd Cosmet. Toxic.*, 14, 809-810.

SCENIHR, 2010

Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf

Sigma-Aldrich

Accessed March 2013. Available at: <http://www.sigmaaldrich.com/united-kingdom.html>

SRC, 2013

Syracuse Research Corporation. Interactive PhysProp Database Demo. Record for cyclotene. Available at: <http://www.syrres.com/what-we-do/databaseforms.aspx?id=386>

Toxtree

Estimation of toxic hazard – a decision tree approach. Ideaconsult Ltd. (Version 2.5.0). Accessed March 2013. Available for download at: http://ihcp.jrc.ec.europa.eu/our_labs/computational_toxicology/qsar_tools/toxtree

Vollmuth, T.A., Bennett, M.B., Hoverman, A.M., Christian, M.S., 1990

An evaluation of food flavoring ingredients using an *in vivo* reproductive and developmental toxicity screening test. *Teratology*, 41, 597.

Withers, R.F.J., 1966

The action of some lactones and related compounds on human chromosomes. Mechanisms of Mutation and Inducing Factors, Symposium Proceedings, 1965. Landa Z. (Ed.), p359-364.