


Additive Toxicity Statement

Piperonal

DESCRIPTION

Name	Piperonal
IUPAC name	1,3-Benzodioxole-5-carboxaldehyde
CAS no	120-57-0
EINECS no	204-409-7
Molecular formula	C ₈ H ₆ O ₃

Structural formula



Molecular weight (g/mol)	150.13
--------------------------	--------

CLP HEALTH HAZARD CLASSIFICATION

This ingredient has been classified under Regulation (EC) No 1272/2008 of the European Parliament and of the Council and is in the classification and labelling inventory.

The CLP classifications of this ingredient are not considered applicable for one, or more, of the following reasons:

- the level required to elicit a response in a toxicological study, with a relevant route of exposure, is far higher than that of its use in BAT products
- the toxicological effect identified does not relate to the route of exposure associated with its use in BAT products
- no relevant toxicological study was identified relating to the route of exposure, and there is only limited consumer exposure to this ingredient associated with its use in BAT products

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are:

Additive Toxicity Statement

Piperonal



TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this, our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch *et al*¹¹.

In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴
- Assessment of the irritation potential of Swedish snus ingredients using the Epioral™ tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself. In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

1. Available at: [http://www.bibra-information.co.uk/supported access to our chemical toxicology database TRACE.html](http://www.bibra-information.co.uk/supported%20access%20to%20our%20chemical%20toxicology%20database%20TRACE.html)
2. Available at: <http://toxnet.nlm.nih.gov/index.html>
3. Available at: <http://ccinfoweb.ccohs.ca/rtecs/search.html>
4. Available at: <http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384>
5. Available at: <http://www.inchem.org/>
6. at: <http://echa.europa.eu/information-on-chemicals>
7. Available at: <http://www.accessdata.fda.gov/scripts/cfn/fcnNavigation.cfm?rpt=eafusListing>
8. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp>
9. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
10. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
11. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem. Toxicol.* 42 Suppl:S53-S83, 2004.
12. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids. *Regul. Toxicol. Pharmacol.* 72 (2):361-369, 2015.
13. B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. *Toxicology* Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
14. Available Letters 229S:S109, 2014. (Abstract)
15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. *The Toxicologist*, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
16. World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004
17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf.
18. SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_051.pdf.

REFERENCES

Adams, T.B., Cohen, S.M., Doull, J., Feron, V.J., Goodman, J.I., Marnett, L.J., Munro, I.C., Portoghese, P.S., Smith, R.L., Waddell, W.J., Wagner, B.M; Expert Panel of the Flavor and Extract Manufacturers Association, 2005
The FEMA GRAS assessment of hydroxy- and alkoxy-substituted benzyl derivatives used as flavor ingredients. *Fd Chem. Toxic.*, 43, 1241-1271.

Alfa Aesar

Accessed March 2013. Available at: <http://www.alfa.com/en/go160w.pgm?srchtyp=product>

Burdock, G.A., 2010

Fenaroli's Handbook of Flavor Ingredients. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.

CCRIS, 2009

Chemical Carcinogenesis Research Information System: piperonal. CCRIS record no. 5928. Last revision date 1 October 2009. Available at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>

ChemIDplus

Accessed April 2013. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>

ChemSpider

Royal Society of Chemistry chemical structure database. Accessed April 2013. Available at: <http://www.chemspider.com/>

CoE, 2000

Chemically-defined flavouring substances. Council of Europe Publishing. ISBN 92-871-4453-2.

ECHA

European Chemicals Agency. Information on Chemicals. Accessed April 2013. Available at: <http://echa.europa.eu/information-on-chemicals>

EFSA, 2012

European Food Safety Authority. Scientific Opinion on Flavouring Group Evaluation 20, Revision 4 (FGE.20Rev4): Benzyl alcohols, benzaldehydes, a related acetal, benzoic acids, and related esters from chemical groups 23 and 30. EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids. *EFSA J.*, 10(12), 2994. Available at: <http://www.efsa.europa.eu/en/search/doc/2994.pdf>

EFSA, 2008

European Food Safety Authority. Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in contact with Food (AFC) on a request from the Commission related to Flavouring Group Evaluation 52 (FGE.52): Consideration of hydroxy- and alkoxy-substituted benzyl derivatives evaluated by JECFA (57th meeting) structurally related to benzyl alcohols, benzaldehydes, a related acetal, benzoic acids, and related esters evaluated by EFSA in FGE.20 (2005) (Commission Regulation (EC) No. 1565/2000 of 18 July 2000). (Question No. EFSA-Q-2008-032C). (Adopted on 3 July 2007). Available at: <http://www.efsa.europa.eu/fr/scdocs/doc/637.pdf>

Additive Toxicity Statement

Piperonal



EFSA, 2012

European Food Safety Authority. EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids (CEF); Scientific opinion on Flavouring Group Evaluation 20, Revision 4 (FGE.20Rev4): Benzyl alcohols, benzaldehydes, a related acetal, benzoic acids, and related esters from chemical groups 23 and 30. Available at: <http://www.efsa.europa.eu/en/efsajournal/doc/2994.pdf>

EPA

US Environmental Protection Agency. Integrated Risk Information System (IRIS). Accessed December 2014. Available at: http://www.epa.gov/iris/search_human.htm and http://cfpub.epa.gov/ncea/iris_drafts/erd.cfm?excCol=Archive&archiveStatus=both

EPI Suite

Accessed March 2013. The database is available to download at: <http://www.epa.gov/opptintr/exposure/pubs/episuitedi.htm>

Epstein, S.S., Arnold, E., Andrea, J., Bass, W., Bishop, Y., 1972

Detection of chemical mutagens by the dominant lethal assay in the mouse. Toxic. appl. Pharmac., 23, 288-325.

FDA, 2011

US Food and Drug Administration. Everything added to Food in the United States (EAFUS). Last updated 17 November 2011. Accessed April 2013. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>

Fisher Scientific, 2011

Material safety datasheet: Piperonal, last revised 10th February 2011. Available at: https://extranet.fisher.co.uk/chemicalProductData_uk/weracs?itemCode=13134-5000

Flavis

EUROPA food flavouring website. Accessed March 2013. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm

Frosch, P.J., Pilz, B., Andersen, K.E., Burrows, D., Camarasa, J.G., Dooms-Goossens, A., Ducombs, G., Fuchs, T., Hannuksela, M., Lachapelle, J.M., Lahti A, Maibach, H.I., Menné T, Rycroft, R.J.G., Shaw, S., Wahlberg, J.E., White, I.R., Wilkinson, J.D., 1995

Patch testing with fragrances: results of a multicenter study of the European Environmental and Contact Dermatitis Research Group with 48 frequently used constituents of perfumes. Contact Dermatitis, 33, 333-342.

Frosch, P.J., Johansen, J.D., Menné, T., Pirker, C., Rastogi, S.C., Andersen, K.E., Bruze, M., Goossens, A., Lepoittevin, J.P., White, I.R., 2002

Further important sensitizers in patients sensitive to fragrances. II. Reactivity to essential oils. Contact Dermatitis, 47, 279-287.

GESTIS

Gefahrstoffinformationssystem (Databases on hazardous substances). Accessed March 2013. Available at: http://limitvalue.ifa.dguv.de/Webform_gw.aspx

Additive Toxicity Statement

Piperonal



Good Scents Company

Accessed March 2013. Available at: <http://www.thegoodscentcompany.com/>

Haworth, S., Lawlor, T., Mortelmans, K., Speck, W., Zeiger, E., 1983

Salmonella mutagenicity test results for 250 chemicals. *Envir. Mutagen.*, 5 (Suppl. 1), 3-49.

HSDB, 2008

Hazardous Substances Data Bank. Last revision date 22 April 2008. Available at:

<http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>

IARC

International Agency for Research on Cancer. World Health Organization. Accessed April 2013.

Available at: <http://www.iarc.fr/>

JECFA, 1967

Toxicological evaluation of some flavouring substances and non-nutritive sweetening agents.

FAO Nutrition Meetings Report Series 44A. WHO/Food Add./68.33. Eleventh Report of the Joint FAO/WHO Expert Committee on Food Additives. Food and Agriculture Organization of the United Nations, World Health Organization, Geneva. Available at:

<http://www.inchem.org/documents/jecfa/jecmono/v44aje32.htm>.

JECFA, 1968

Specifications for the identity and purity of food additives and their toxicological evaluation:

some flavouring substances and non-nutritive sweetening agents. WHO Technical Report Series 383. FAO Nutrition Meetings Report Series 44. Eleventh Report of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization, Geneva. Available at:

http://whqlibdoc.who.int/trs/WHO_TRS_383.pdf.

JECFA, 2002a

Evaluation of certain food additives and contaminants. WHO Technical Report Series 909. Fifty-seventh report of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization, Geneva. Available at: http://whqlibdoc.who.int/trs/WHO_TRS_909.pdf.

JECFA, 2002b

Safety evaluation of certain food additives and contaminants. Hydroxy- and alkoxy-substituted benzyl derivatives. WHO Food Additives Series 48. Prepared by the fifty-seventh meeting of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization, Geneva.

Available at: <http://www.inchem.org/documents/jecfa/jecmono/v48je15.htm>.

JRC

Joint Research Centre. European chemical Substances Information System (ESIS). Accessed March 2013. Available at: <http://esis.jrc.ec.europa.eu/>

Kasamaki, A. and Urasawa, S., 1985

Transforming potency of flavoring agents in Chinese hamster cells. *J. Toxic. Sci.*, 10, 177-185.

Kasamaki, A., Takahashi, H., Tsumura, N., Niwa, J., Fujita, T., Urasawa, S., 1982

Genotoxicity of flavoring agents. *Mutation Res.*, 105, 387-392.

Katz, A.E., 1946

Dermal irritating properties of essential oils and aromatic chemicals. *Spice Mill*, 69, 46-51.

Klecak, G., Geleick, H., Frey, J.R., 1977

Screening of fragrance materials for allergenicity in the guinea-pig. I. Comparison of four testing methods. *J. Soc. cosmet. Chem.*, 28, 53-64.

Lorber, M., 1972

Hematotoxicity of synergized pyrethrin insecticides and related chemicals in intact, totally and subtotally splenectomized dogs. *Acta hepato-Gastroent.*, 19, 66-78.

Merck, 2006

The Merck Index. An Encyclopedia of Chemicals, Drugs and Biologicals. Ed. O'Neil J et al, Fourteenth edition. Merck & Co., Ltd., Whitehouse Station, NJ, USA.

NTP

US National Toxicology Program. Accessed December 2014. Available at:
<http://www.niehs.nih.gov/research/resources/databases/cebs/index.cfm> and
<http://tools.niehs.nih.gov/cebs3/ui/>

NTP, 2000

National Toxicology Program. Salmonella Ames test: piperonal. Study ID A18234. Available at:
http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm?fuseaction=salmonella.salmonellaData&study_no=A18234&cas_no=120%2D57%2D0&endpointlist=SA

NTP, 1980

National Toxicology Program. Salmonella Ames test: piperonal. Study ID 287791. Available at:
http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm?fuseaction=salmonella.salmonellaData&study_no=287791&cas_no=120%2D57%2D0&endpointlist=SA

Oda, Y., Hamono, Y., Inoue, K., Yamamoto, H., Niihara, T., Kunita, N., 1978

Mutagenicity of food flavours in bacteria (1st report). *Osaka Furitsu KEKHSEHI*, 9, 177-181 (in Japanese, tables in English).

Opdyke, D.L.J., 1974

Monographs on fragrance raw materials: Heliotropin. *Fd Cosmet. Toxicol.*, 12, 907-908.

RTECS, 2003

Registry of Toxic Effects of Chemical Substances. Record for Piperonal (120-57-0). Last updated November 2003.

SCENIHR, 2010

Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at:
http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf

Additive Toxicity Statement

Piperonal



Sekizawa, J. and Shibamoto, T., 1982

Genotoxicity of safrole-related chemicals in microbial test systems. *Mutation Res.*, 101, 127-140.

Sigma-Aldrich, 2012

Material safety datasheet: Piperonal, last revised 18 September 2012. Available at:

<http://www.sigmaaldrich.com/MSDS/MSDS/DisplayMSDSPage.do?country=GB&language=en&productNumber=P49104&brand=ALDRICH&PageToGoToURL=http%3A%2F%2Fwww.sigmaaldrich.com%2Fcatalog%2Fproduct%2Faldrich%2Fp49104%3Flang%3Den>

SRC, 2013

Syracuse Research Corporation. Interactive PhysProp Database Demo. Available at:

<http://www.syrres.com/what-we-do/databaseforms.aspx?id=386>

ToxTree

Estimation of toxic hazard – a decision tree approach. Version 2.6.0. IdeaConsult Ltd. Available at: <http://toxtree.sourceforge.net>.

White, T.J., Goodman, D., Shulgin, A.T., Castagnoli Jr, N., Lee, R., Petrakis, N.L., 1977

Mutagenic activity of some centrally active aromatic amines in *Salmonella typhimurium*. *Mutation Res.*, 56, 199-202.

Additive Toxicity Statement

Piperonal

