

Additive Toxicity Statement

Titanium dioxide



DESCRIPTION

Name	Titanium dioxide
IUPAC name	Dioxotitanium
CAS no	13463-67-7
EINECS no.	236-675-5
Molecular formula	TiO ₂
Structural formula	



Molecular weight (g/mol)	79.865
--------------------------	--------

AVAILABLE STUDIES

The available toxicity studies indicated in the Commission Implementing Decision (EU) 2015/2186 have been provided as a bibliography of published papers. The studies were obtained by BAT for toxicological assessments, to ensure that additives do not increase the inherent risk associated with the use of our products.

The risk assessment starts with a comprehensive search for relevant papers, using the additive's name, major synonyms and CAS Registry Number. The main sources searched are: TRACE¹, Toxnet², RTECS³, TSCATS⁴, INCHEM⁵, Europa Food Flavouring⁶, ECHA⁷, EAFUS⁸, ChemIDplus⁹ and eChemportal¹⁰.

RISK ASSESSMENT

Toxicological assessments are carried out by our scientists (including a number of European Registered Toxicologists (ERT)) at our Research and Development facilities in the UK. Our approach excludes the use of formally classified genotoxicants, non-threshold carcinogens, mutagens, reproductive and developmental toxicants as additives. Based on Levels of Concern and weight-of evidence, our approach ensures that additives are used at levels lower than the relevant toxicological reference value.

Following a comprehensive search for all available toxicological information, our toxicologists select the most appropriate studies for evaluation for the intended route of exposure. To do this, our toxicologists evaluate the quality of all pertinent studies identified and the data used. The evaluation of data quality includes an assessment of its relevance and reliability as well as the adequacy of the information for hazard/risk assessment purposes, following the principles described by Klimisch *et al*¹¹.

Additive Toxicity Statement

Titanium dioxide



In the majority of BAT's products, a number of the additives are heated or combusted. The effects of heating or combustion on additive toxicity, have been addressed by extensive testing. The results of pyrolysis, smoke chemistry, *in vitro* cytotoxicity, *in vitro* genotoxicity, inhalation toxicity and tumourigenicity studies have been widely published in peer-reviewed journals. These studies are included in our risk assessments where applicable by product class.

Examples of our assessment processes can be found in published literature for example:

- An overview of the effects of tobacco ingredients on smoke chemistry and toxicity¹²
- An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids¹³
- Contact sensitisation risk assessment approach for pouched snus ingredients¹⁴
- Assessment of the irritation potential of Swedish snus ingredients using the Epioral™ tissue model¹⁵

Further examples of our scientific publications are available at www.bat-science.com.

Health risks of tobacco use have primarily been determined in long term human epidemiological studies. For example, the smoking population in countries such as Canada, Australia and the UK have historically smoked Virginia style cigarettes, which contain few additives. In other countries such as the US and Germany smokers prefer American-blended style cigarettes, which contain significantly more additives. Notwithstanding the distinction in historical use of additives in these countries, there appears to be no obvious difference in the relative risks of cigarette smoking for these types of cigarette, or on the incidence of diseases such as lung cancer and chronic obstructive pulmonary disease¹⁶, suggesting that the addition of additives to cigarettes may not increase the incidence of diseases associated with smoking.

ADDICTIVENESS

In its 2010 opinion on Addictiveness and Attractiveness of additives¹⁷, SCENIHR came to the clear conclusion that no additive could be identified which has an "addictive" effect in isolation, and that there are no indications that additives increase the "addictive" effect of nicotine itself. In a more recent final opinion¹⁸, SCENIHR reviewed 1260 additives and selected only 14 substances for further study because of their contribution to addictiveness to smoking.

CONCLUSION

Based on the available scientific evidence, BAT's scientists have concluded that the additives used in BAT's tobacco products, do not add to the toxicological risks of using those products.

1. Available at: <http://www.bibra-information.co.uk/supported> access to our chemical toxicology database TRACE.html
2. Available at: <http://toxnet.nlm.nih.gov/index.html>
3. Available at: <http://ccinfoweb.ccohs.ca/rtecs/search.html>
4. Available at: <http://www.srcinc.com/what-we-do/databaseforms.aspx?id=384>
5. Available at: <http://www.inchem.org/>
6. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm
7. Available at: <http://echa.europa.eu/information-on-chemicals>
8. Available at: <http://www.accessdata.fda.gov/scripts/cfn/fcnNavigation.cfm?rpt=eafusListing>
9. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp>

Additive Toxicity Statement

Titanium dioxide



10. Available at: http://www.echemportal.org/echemportal/index?pageID=0&request_locale=en
 11. Klimisch, H.J., Andreae, E., Tillmann, U., (1997). A systematic approach for evaluating the quality of experimental and ecotoxicological data. *Regul. Toxicol. Pharmacol.* 25, 1–5.
 12. R. R. Baker, E. D. Massey and G. Smith. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem. Toxicol.* 42 Suppl:S53-S83, 2004.
 13. S. Costigan and C. Meredith. An approach to ingredient screening and toxicological risk assessment of flavours in e-liquids. *Regul. Toxicol. Pharmacol.* 72 (2):361-369, 2015.
 14. B. Lang, S. Costigan, S. Goodall and C. Meredith. Contact sensitisation risk assessment approach for pouched snus ingredients. *Toxicology Letters* 229S:S109, 2014. (Abstract)
 15. L. Neilson, S. Faux, S., Hinchcliffe, T. Jai and C. Meredith. Assessment of the irritation potential of swedish snus ingredients using the epioral™ tissue model. Society of Toxicology, Baltimore, USA, March 15-19th. *The Toxicologist*, Volume 108, no 1, pg 307-308 (March 2009) (Conference Poster)
 16. World Health Organisation, 2004. Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 83. Tobacco smoke and involuntary smoking. p 171. International Agency for Research on Cancer (IARC), Lyon, 2004
 17. SCENIHR, 2010. Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. ISBN 978-92-79-12788-5. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf.
 18. SCENIHR, 2015. Final Opinion on Additives used in Tobacco Products (Opinion 1). The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_051.pdf.
-

REFERENCES

Burdock, G.A., 2010

Fenaroli's Handbook of Flavor Ingredients. 6th Edition. CRC Press, Boca Raton. ISBN 978-1-4200-9077-2.

ChemIDplus

Record for Titanium dioxide, CAS RN 13463-67-7. Accessed November 2014. Available at: <http://chem.sis.nlm.nih.gov/chemidplus/rn/13463-67-7>

ChemSpider

Record for Titanium dioxide, CAS RNs covered; 13463-67-7, 1317-80-2, 1317-70-0. Accessed November 2014. Available at: <http://www.chemspider.com/Chemical-Structure.24256.html>

Codex

Codex Alimentarius. International Food Standards. World Health Organization, Food and Agriculture Organization of the United Nations. Last updated 24 January 2014. Available at: <http://www.codexalimentarius.org/codex-home/en/>

CoE, 2000

Chemically-defined flavouring substances. Council of Europe Publishing. ISBN 92-871-4453-2.

CoE, 1992

Flavouring substances and natural sources of flavourings .Volume I. 4th Edition. Chemically-defined flavouring substances. Council of Europe Publishing. ISBN 2-7160-0147-2.

CoE, 1981

Flavouring substances and natural sources of flavourings 3rd Edition. Maisonneuve. ISBN 27160-0081-6.

Dunkel, V.C., Zeiger, E., Brusick, D., McCoy, E., McGregor, D., Mortelmans, K., Rosenkranz, H.S., Simmon, V.F., 1985

Reproducibility of microbial mutagenicity assays. II. Testing of carcinogens and noncarcinogens in *Salmonella typhimurium* and *Escherichia coli*. *Envir. Mutagen.*, 7 (Suppl. 5), 1-248.

ECHA, 2014a

European Chemicals Agency. Information on Chemicals. Dossier on Titanium dioxide, CAS RN 13463-67-7. First published 17 March 2011, Last modified 28 October 2014. Registrants/suppliers Alfarben, S.A. et al. Available at http://apps.echa.europa.eu/registered/data/dossiers/DISS-9eaff323-014a-482f-e044-00144f67d031/DISS-9eaff323-014a-482f-e044-00144f67d031_DISS-9eaff323-014a-482f-e044-00144f67d031.html

ECHA, 2014b

European Chemicals Agency. Classification and Labelling (C&L) Inventory database. Last updated 3 October 2014. Available at: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database>

Additive Toxicity Statement

Titanium dioxide



EFSA

European Food Safety Authority. Accessed November 2014. Available at: <http://www.efsa.europa.eu/>

EPA

US Environmental Protection Agency. Integrated Risk Information System (IRIS). Accessed November 2014. Available at: http://www.epa.gov/iris/search_human.htm and http://cfpub.epa.gov/ncea/iris_drafts/erd.cfm?excCol=Archive&archiveStatus=both

EPISuite

Accessed November 2014. The database is available to download at: <http://www.epa.gov/opptintr/exposure/pubs/episuitedi.htm>

FDA, 2013a

US Food and Drug Administration. Everything added to Food in the United States (EAFUS). Last updated 23 April 2013. Accessed November 2014. Available at: <http://www.accessdata.fda.gov/scripts/fcn/fcnNavigation.cfm?rpt=eafusListing>

FDA, 2013b

US Food and Drug Administration. Food Additive Status List. Last updated 21 March 2013. Available at: <http://www.fda.gov/Food/IngredientsPackagingLabeling/FoodAdditivesIngredients/ucm091048.htm#ftnP>

Flavis

EUROPA food flavouring website. Accessed November 2014. Available at: http://ec.europa.eu/food/food/chemicalsafety/flavouring/database/dsp_search.cfm

GESTIS

Gefahrstoffinformationssystem (Databases on hazardous substances). Accessed November 2014. Available at: http://limitvalue.ifa.dguv.de/Webform_gw.aspx

Good Scents Company

Accessed November 2014. Available at: <http://www.thegoodscentscompany.com/>

HSDB, 2014

Hazardous substance database, record for titanium dioxide, reviewed on 16 January 2014. Available at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search2/f?/.temp/~qjDXRw:1>

IARC, 2010

International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Volume 93. Carbon black, titanium dioxide, and talc. IARC Working Group on the Evaluation of Carcinogenic Risks to Human, Lyon, 7-14 February 2006. Available at: <http://monographs.iarc.fr/ENG/Monographs/vol93/mono93.pdf>

Additive Toxicity Statement

Titanium dioxide



JECFA, 2012

Evaluation of certain food additives. *WHO tech. Rept Ser.*, 974. Seventy-sixth report of the Joint FAO/WHO Expert Committee on Food Additives. Geneva, 5-14 June 2012. Available at: http://apps.who.int/iris/bitstream/10665/77752/1/WHO_TRS_974_eng.pdf?ua=1

JECFA

Joint Expert Committee on Food Additives. Monographs and Evaluations. Accessed November 2014. Available at: <http://www.inchem.org/pages/jecfa.html>

Lu, P-J., Ho, I.C., Lee, T.C., 1998

Induction of sister chromatid exchanges and micronuclei by titanium dioxide in Chinese hamster ovary-K1 cells. *Mutation Res.*, 414, 15-20.

Merck, 2013

The Merck Index. An Encyclopedia of Chemicals, Drugs and Biologicals. Ed. O'Neil MJ et al. Fifteenth edition. Royal Society of Chemistry, Cambridge, UK.

Miller, B.M., Pujadas, E., Gocke, E., 1995

Evaluation of the micronucleus test *in vitro* using Chinese hamster cells: results of four chemicals weakly positive in the *in vivo* micronucleus test. *Envir. molec. Mutagen.*, 26, 240-247.

NCI, 1979

US National Cancer Institute. Bioassay of titanium dioxide for possible carcinogenicity. CAS No. 13463-67-7. NCI-CG-TR97. Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute. Available at: http://ntp.niehs.nih.gov/ntp/htdocs/lt_rpts/tr097.pdf

NSF, 2005

NSF International. Titanium and titanium dioxide: oral risk assessment – executive summary. 10 June 2005. Abstract available at: http://www.techstreet.com/direct/nsf/titanium_exec_sum.pdf

NTP, 2004

US National Toxicology Program. NTP studies on titanium dioxide (CAS RN 13463-67-7). Available at: http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm

NTP, 1989

US National Toxicology Program. NTP study 860815.

NTP

US National Toxicology Program. Accessed November 2014. Available at: <http://www.niehs.nih.gov/research/resources/databases/cebs/index.cfm> and <http://tools.niehs.nih.gov/cebs3/ui/>

SCCNFP, 2000

Opinion of the Scientific Committee on Cosmetic Products and Non-Food Products Intended for Consumers concerning titanium dioxide. Colipa No. S75. SCCNFP/0005/98, adopted by the SCCNFP during the 14th plenary meeting of 24 October 2000. Available at: http://ec.europa.eu/health/ph_risk/committees/sccp/documents/out135_en.pdf

Additive Toxicity Statement

Titanium dioxide



SCENIHR, 2010

Addictiveness and Attractiveness of Tobacco Additives. The Scientific Committee on Emerging and Newly Identified Health Risks. European Union. ISBN 978-92-79-12788-5. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_029.pdf

Sheftel, V.O., 1990

Toxic Properties of Polymers and Additives. Inglis E and Dunstall S (Eds). Rapra Technology Limited.

SRC, 2013

Syracuse Research Corporation. Interactive PhysProp Database Demo. Available at: <http://esc.syrres.com/fatepointer/search.asp>

Toxtree

Estimation of toxic hazard – a decision tree approach. Ideaconsult Ltd. (Version 2.6.6). Accessed November 2014. Available for download at: http://ihcp.jrc.ec.europa.eu/our_labs/computational_toxicology/qsar_tools/toxtree

Warheit, D.B., Brock, W.J., Lee, K.P., Webb, T.R., Reed, K.L., 2005

Comparative pulmonary toxicity inhalation and instillation studies with different TiO₂ particle formulations: impact of surface treatments on particle toxicity. *Toxic. Sci.*, 514-524.

Wilfong, E.R., Lyles, M., Rietcheck, R.L., Arfsten, D.P., Boeckman, H.J., Johnson, E.W., Doyle, T.L., Chapman, G.D., 2011

The acute and long-term effects of Middle East sand particles on the rat airway following a single intratracheal instillation. *J. Toxicol. envir. Hlth Part A*, 74, 1351-1365.