

Overview information for

Benzo-(alpha) pyrene

BENZO-(ALPHA) PYRENE REFERENCES					
Author Name	Title	Journal	Volume	Page number(s)	Year
Li X, Luo Y, Jiang X, Zhang H, Zhu F, Hu S, Hou H, Hu Q, Pang Y.	Chemical Analysis and Simulated Pyrolysis of Tobacco Heating System 2.2 Compared to Conventional Cigarettes.	Nicotine Tob Res.	21(1)	111-118	2019
Lorkiewicz P, Riggs DW, Keith RJ, Conklin DJ, Xie Z, Sutaria S, Lynch B, Srivastava S, Bhatnagar A.	Comparison of Urinary Biomarkers of Exposure in Humans Using Electronic Cigarettes, Combustible Cigarettes, and Smokeless Tobacco.	Nicotine Tob Res.	21(9)	1228-1238	2019
Cai B, Li Z, Wang R, Geng Z, Shi Y, Xie S, Wang Z, Yang Z, Ren X.	Emission level of seven mainstream smoke toxicants from cigarette with variable tobacco leaf constituents.	Regul Toxicol Pharmacol.	103	181-188	2019
Weng MW, Lee HW, Park SH, Hu Y, Wang HT, Chen LC, Rom WN, Huang WC, Lepor H, Wu XR, Yang CS, Tang MS.	Aldehydes are the predominant forces inducing DNA damage and inhibiting DNA repair in tobacco smoke carcinogenesis.	Proc Natl Acad Sci U S A.	115(27)	E6152-E6161	2018
Zhang Y, Zou HY, Shi P, Yang Q, Tang LJ, Jiang JH, Wu HL, Yu RQ.	Determination of benzo[a]pyrene in cigarette mainstream smoke by using mid-infrared spectroscopy associated with a novel chemometric algorithm.	Anal Chim Acta.	902	Sep-43	2016
Thorne D, Crooks I, Hollings M, Seymour A, Meredith C, Gaca M.	The mutagenic assessment of an electronic-cigarette and reference cigarette smoke using the Ames assay in strains TA98 and TA100.	Mutat Res.	812	29-38	2016

Yershova K, Yuan JM, Wang R, Valentin L, Watson C, Gao YT, Hecht SS, Stepanov I.	Tobacco-specific N-nitrosamines and polycyclic aromatic hydrocarbons in cigarettes smoked by the participants of the Shanghai Cohort Study.	Int J Cancer.	139(6)	1261-9	2016
Wang H, Li X, Guo J, Peng B, Cui H, Liu K, Wang S, Qin Y, Sun P, Zhao L, Xie F, Liu H.	Distribution of toxic chemicals in particles of various sizes from mainstream cigarette smoke.	Inhal Toxicol.	28(2)	89-94	2016
Vu AT, Taylor KM, Holman MR, Ding YS, Hearn B, Watson CH.	Polycyclic Aromatic Hydrocarbons in the Mainstream Smoke of Popular U.S. Cigarettes.	Chem Res Toxicol.	28(8)	1616-26	2015
Machado Jde B, Chatkin JM, Zimmer AR, Goulart AP, Thiesen FV.	Cotinine and polycyclic aromatic hydrocarbons levels in the amniotic fluid and fetal cord at birth and in the urine from pregnant smokers.	PLoS One.	9(12):e116293.		2014
Ding YS, Ward J, Hammond D, Watson CH.	Mouth-level intake of benzo[a]pyrene from reduced nicotine cigarettes.	Int J Environ Res Public Health	11(11)	11898-914	2014
Zhang X, Hou H, Shi L, Liu Y, Wang A, Hu Q.	Novel method to analysis benzo[a]pyrene in filter by liquid chromatography/tandem mass spectrometry: application to assess mouth level benzo[a]pyrene exposure.	Rapid Commun Mass Spectrom.	28(13)	1468-72	2014
Dittrich David J; Fieblekorn Richard T; Bevan Michael J; Rushforth David; Murphy James J; Ashley Madeleine; McAdam Kevin G; Liu Chuan; Proctor Christopher J	Approaches for the design of reduced toxicant emission cigarettes.	SpringerPlus	3	374	2014

Slotkin Theodore A; Card Jennifer; Stadler Ashley; Seidler Frederic J; Levin Edward D	Effects of tobacco smoke on PC12 cell neurodifferentiation are distinct from those of nicotine or benzo[a]pyrene.	Neurotoxicology and teratology	43	19-24	2014
Siddique Shabana; Sadeu Jean C; Foster Warren G; Feng Yong- Lai; Zhu Jiping	In vitro exposure to cigarette smoke induces oxidative stress in follicular cells of F1 hybrid mice.	Journal of applied toxicology	34	224-6	2014
Hyodo T; Minagawa K; Inoue T; Fujimoto J; Minami N; Bito R; Mikita A	Estimation of mouth level exposure to smoke constituents of cigarettes with different tar levels using filter analysis.	Regulatory toxicology and pharmacology	67	486-98	2013
Garcia-Canton Carolina; Anadon Arturo; Meredith Clive	Genotoxicity evaluation of individual cigarette smoke toxicants using the in vitro γ H2AX assay by high content screening.	Toxicology letters	223	Jul-81	2013
Yamamoto Mitsuko L; Chapman Aaron M; Schiestl Robert H	Effects of side-stream tobacco smoke and smoke extract on glutathione-and oxidative DNA damage repair-deficient mice and blood cells.	Mutation research	749	58-65	2013
Sadeu Jean Clair; Foster Warren G	The cigarette smoke constituent benzo[a]pyrene disrupts metabolic enzyme, and apoptosis pathway member gene expression in ovarian follicles.	Reproductive toxicology	40	Sep-52	2013
Jaiswal Aruna S; Panda Harekrushna; Pampo Christine A; Siemann Dietmar W; Gairola C Gary; Hromas Robert; Narayan Satya	Adenomatous polyposis coli-mediated accumulation of abasic DNA lesions lead to cigarette smoke condensate-induced neoplastic transformation of normal breast epithelial cells.	Neoplasia	15	454-60	2013

Piade J-J; Wajrock S; Jaccard G; Janeke G	Formation of mainstream cigarette smoke constituents prioritized by the World Health Organization--yield patterns observed in market surveys, clustering and inverse correlations.	Food and chemical toxicology	55	329-47	2013
Ashraf, M. Ejaz, S. Kim, B. Lim, C.W. Nawaz, M.	Cigarette smoke condensate and total particulate matter severely disrupts physiological angiogenesis	Food and Chemical Toxicology	47(3)	601-614	2009
Ashraf, M. Ejaz, S. Kim, B. Lim, C.W. Nawaz, M.	Anti-angiogenic and teratological activities associated with exposure to total particulate matter from commercial cigarettes	Food and Chemical Toxicology	47(2)	368-376	2009
Berger, A. Hamm, J.T. Misra, M. Mitchell, M. Vulimiri, S.V.	Effects of Mainstream Cigarette Smoke on the Global Metabolome of Human Lung Epithelial Cells	Chem. Res. Toxicol.	22	492-503	2009
Berndt-Weis, L. Douglas, G.R. Halappanavar, S. Stampfli, M.R. Williams, A. Yauk, C.L.	Toxicogenomic Analysis of Mainstream Tobacco Smoke-Exposed Mice Reveals Repression of Plasminogen Activator Inhibitor-1 Gene in Heart	Inhalation Toxicology	21(1-4)	78-85	2009
Bhattacharya, A. Chakrabarti, G. Das, A.	Cigarette Smoke Extract Induces Disruption of Structure and Function of Tubulin-Microtubule in Lung Epithelium Cells and in Vitro	Chem. Res. Toxicol.	22	446-459	2009
Chen, C.-C. Chen, R.-M. Pan, W.-C. Shen, Y.-C. Ueng, Y.-F.	Suppressive effect of tobacco smoke extracts on oral P-glycoprotein function and its impact in smoke-induced insult to oral epidermal cells	Toxicology Letters	185	116-123	2009
Chen, C.-C. Chen, R.-M. Pan, W.-C. Shen, Y.-C. Ueng, Y.-F.	Suppressive effect of tobacco smoke extracts on oral P-glycoprotein function and its impact in smoke-induced insult to oral epidermal cells	Toxicology Letters	185	116-123	2009

Chen, H.-W. Ku, H.-J. Lii, C.-K. Wang, T.-S.	Cigarette Smoke Extract Induces Expression of Cell Adhesion Molecules in HUVEC via Actin Filament Reorganization	Environmental and Molecular Mutagenesis	50	96-104	2009
Kim, I.k. Kim, K.-N. Kim, M.-K. Lee, J. Sohn, S.-H.	Effect of tobacco compounds on gene expression profiles in human epithelial cells	Environmental Toxicology and Pharmacology	27	111-119	2009
Kumaravel, T.S. Massey, E.D. McEwan, M. Thorne, D. Wilson, J.	Measurement of oxidative DNA damage induced by mainstream cigarette smoke in cultured NCI-H292 human pulmonary carcinoma cells	Mutation Research	673	08-Mar	2009
Al-Mugotir, M. Bitterman, P. Fang, Q. Kim, H. Kobayashi, T. Liu, X. Mao, L. Rennard, S. Togo, S. Wang, X.	NF-kappaB mediates the survival of human bronchial epithelial cells exposed to cigarette smoke extract	Respiratory Research	9(66)	11-Jan	2008
Aufderheide, M. Gressmann, H.	Mutagenicity of native cigarette mainstream smoke and its gas/vapour phase by use of different tester strains and cigarettes in a modified Ames assay	Mutation Research	656	82-87	2008
Carter, J.D. Crissman, K. Dailey, L.A. Foronjy, R.F. Ghio, A.J. Hilborn, E.D. Pinkerton, K.E. Richards, J.H. Stonehuerner, J. Uyeminami, D.	Particulate Matter in Cigarette Smoke Alters Iron Homeostasis to Produce a Biological Effect	Am J Respir Crit Care Med	178	1130-1138	2008

Casola, A. Castro, S.M. Garofalo, R.P. Guerrero-Plata, A. Kolli, D.	Cigarette Smoke Condensate Enhances Respiratory Syncytial Virus-Induced Chemokine Release by Modulating NF-kappa B and Interferon Regulatory Factor Activation	Toxicological Sciences	106(2)	509-518	2008
Coggins, C.R.E. Friedrichs, B. Gerstenberg, B. Gomm, W. Meurrens, K. Patskan, G.J. Podraza, K.F. Schnell, P. Stabbert, R. Terpstra, P. Veltel, D. Weber, S.	Toxicological Comparisons of Three Styles of a Commercial U.S. Cigarette (Marlboro) with the 1 R4F Reference Cigarette	Inhalation Toxicology	20(5-8)	695-721	2008
DeMarini, D.M. Gudi, R. Kehl, M. Kirby, P.E. Polzin, G. Rao, M. Recio, L. Richter, P.A. Szkudlinska, A.	Genotoxicity of 10 cigarette smoke condensates in four test systems: Comparisons between assays and condensates	Mutation Research	650	15-29	2008
Edmiston, J. Flora, J.W. Langston, T.B. Li, G. McKinney, W. Rana, G.S.J.B. Secrist, R.	Identification of in vitro differential cell secretions due to cigarette smoke condensate exposure using nanoflow capillary liquid chromatography and high-resolution mass spectrometry	Anal Bioanal Chem	391	2845-2856	2008
Edmiston, J.S. Kane, D.B. McKinney, W.J. Oldham, M.J. Scian, M.J.	Characterization of a Whole Smoke In Vitro Exposure System (Burghart Mimic Smoker-01)	Inhalation Toxicology	XX	10-Jan	2008
Edvinsson, L. Xu, C.-B. Zhang, W. Zhang, Y. Zheng, J.-P.	Lipid-Soluble Smoke Particles Upregulate Vascular Smooth Muscle ETB Receptors via Activation of Mitogen-Activating Protein Kinases and NF- kappaB Pathways	Toxicological Sciences	106(2)	546-555	2008

Kaushik, G. Kaushik, T. Khanduja, K.L. Khanduja, S. Pathak, C.M.	Cigarette smoke condensate promotes cell proliferation through disturbance in cellular redox homeostasis of transformed lung epithelial type-II cells	Cancer Letters	270	120-131	2008
Klus, H. Muller, L. Thielen, A.	Tobacco smoke: Unraveling a controversial subject	Experimental and Toxicologic Pathology	60(2-3)	141-156	2008
Ambrose, V. Dickson, S.J. Fowles, J. Hampton, S. Lea, R.A. Miller, J.H. Truman, P.	Tobacco particulate matter is more potent than nicotine at upregulating nicotinic receptors on SH-SY5Y cells	Nicotine & Tobacco Research	9(8)	793-799	2007
Aufderheide, M. Gressmann, H.	A modified Ames assay reveals the mutagenicity of native cigarette mainstream smoke and its gas vapour phase	Experimental and Toxicologic Pathology	58	383-392	2007
Avunduk, M.C. Civelek, S. Guyen, C. Isik, A.C.U. Yardimci, S.	Morphologic Alteration Induced by Short-Term Smoke Exposure in Rats	ORL	69	13-17	2007
Baty, C.J. Choi, A.M.K. Guo, F. Karlsson, J.M. Kauffman, H.F. Kim, H.P. Lee, J.S. Liu, F. Postma, D.S. Ryter, S.W. Slebos, D.-J. van der Toorn, M. Wang, X. Watkins, S.C.	Mitochondrial Localization and Function of Heme Oxygenase-1 in Cigarette Smoke-Induced Cell Death	Am J Respir Cell Mol Biol	36	409-417	2007
Bergren, D.R.	Tobacco Smoke is an Adjuvant for Maintained Airway Sensitization in Guinea Pigs	Journal of Asthma	44	723-728	2007
Choi, A.M.K. Dong, Y. Feghali-Bostwick, C.A. Li, C. Matthay, M.A. Ning, W. Sun, J.	Cigarette Smoke Stimulates Matrix Metalloproteinase-2 Activity via EGR-1 in Human Lung Fibroblasts	Am J Respir Cell Mol Biol	36	480-490	2007

Churg, A. Kerstjens, H.A.M. Postma, D.S. Timens, W. Whittaker, P. Wright, J.L.	Airway Remodeling in the Smoke Exposed Guinea Pig Model	Inhalation Toxicology	19	915-923	2007
Coggins, C.R.E.	An Updated Review of Inhalation Studies with Cigarette Smoke in Laboratory Animals	International Journal of Toxicology	26	331-338	2007
Geerlings, M. Hylkema, M.N. Kerstjens, H.A. Luinge, M.A. Melgert, B.N. Postma, D.S. Schouten, J.P. Timens, W.	Effects of 4 months of smoking in mice with ovalbumin-induced airway inflammation	Clinical and Experimental Allergy	37	1798-1808	2007
Hayes, J.R. Meckley, D.R. Nelson, P.R. Stavanja, M.S. Swauger, J.E. Van Kampen, K.R.	Effect of a flue-curing process that reduces tobacco specific nitrosamines on the tumor promotion in SENCAR mice by cigarette smoke condensate	Food and Chemical Toxicology	45	419-430	2007
Hitchman, K. Kerepesi, L. Lu, B. Meng, Q.R. Wisse, L.	Cytotoxicity and Gene Expression Profiles in Cell Cultures Exposed to Whole Smoke from Three Types of Cigarettes	Toxicological Sciences	98(2)	469-478	2007
Ho, C.-Y. Lan, M.-Y. Lee, T.-C. Yang, A.-H.	Cigarette smoke extract induces cytotoxicity on human nasal epithelial cells	Am J Rhinol	21	218-223	2007
Hu, J. Tian, F. Xu, Y.-j. Zhang, Z.-x.	Effect of cigarette smoke extract on proliferation of rat pulmonary artery smooth muscle cells and the relevant roles of protein kinase C	Chin Med J	120(17)	1523-1528	2007
Iann, M.C. Johnson, M. Martin, M.B. Reiter, R. Richards, J.K. Shah, M.S. Singh, B. Stoica, A. Wang, A.	Effects of Tobacco Smoke Condensate on Estrogen Receptor- α Gene Expression and Activity	Endocrinology	148(10)	4676-4686	2007

Lauterbach, J.H. Momin, R.A. Rickert, W.S. Trivedi, A.H. Wright, W.G.	Effect of Smoking Conditions and Methods of Collection on the Mutagenicity and Cytotoxicity of Cigarette Mainstream Smoke	Toxicological Sciences	96(2)	285-293	2007
Meurrens K, Ruf S, Ross G, Schleef R, von Holt K, Schluter K-D	Smoking accelerates the progression of hypertension-induced myocardial hypertrophy to heart failure in spontaneously hypertensive rats.	Cardiovascular Research	76	311-322	2007
Ayres, P.H. Curtin, G.M. Meckley, D.R. Nelson, P.R. Stavanja, M.S. Swauger, J.E.	DBA/2 mouse skin is unresponsive to dermal tumor promotion by cigarette smoke condensate	Experimental and Toxicologic Pathology	58	125-132	2006
Baginski, T.K. Dabbagh, K. Satjawatcharaphong, C. Swinney, D.C.	Cigarette Smoke Synergistically Enhances Respiratory Mucin Induction by Proinflammatory Stimuli	Am J Respir Cell Mol Biol	35	165-174	2006
Bauter, M.R. Chida, A.S. Kilty, I. Maggirwar, S.B. Rahman, I. Seweryniak, K. Shafiq, N. Yang, S.-R.	Cigarette smoke induces proinflammatory cytokine release by activation of NF-kB and posttranslational modifications of histone deacetylase in macrophages	Am J Physiol Lung Cell Mol Physiol	291	46-57	2006
Borlak, J. Gebel, S. Gerstmayer, B. Kuhl, P. Meurrens, K. Muller, T.	The Kinetics of Transcriptomic Changes Induced by Cigarette Smoke in Rat Lungs Reveals a Specific Program of Defense, Inflammation, and Circadian Clock Gene Expression	Toxicological Sciences	93(2)	422-431	2006
Brys, A.M. Gideon, K.M. Harbo, S.J. Jones, R. Lee, M.K. Meng, Q.R. Renne, R.A.	Gene Expression Profiling in Lung Tissues from Mice Exposed to Cigarette Smoke, Lipopolysaccharide, or Smoke Plus Lipopolysaccharide by Inhalation	Inhalation Toxicology	18	555-568	2006

Diekmann, J. Muller, B.L. Tewes, F.J. Wooten, J.B.	Involvement of Semiquinone Radicals in the in Vitro Cytotoxicity of Cigarette Mainstream Smoke	Chem. Res. Toxicol.	19	1602-1610	2006
Espiritu, I. Maronpot, R.R. Witschi, H.	Lung tumors in 2 year old strain A/J mice exposed for 6 months to tobacco smoke	Cancer Letters	241	64-68	2006
Feng, Z. Hu, W. Hu, Y. Tang, M.-s.	Acrolein is a major cigarette-related lung cancer agent: Preferential binding at p53 mutational hotspots and inhibition of DNA repair	PNAS (Proceedings of the National Academy of Sciences)	103(42)	15404-15409	2006
Friedrichs, B. Van Miert, E. Vanscheeuwijck, P.	LUNG INFLAMMATION IN RATS FOLLOWING SUBCHRONIC EXPOSURE TO CIGARETTE MAINSTREAM SMOKE	Experimental Lung Research	32	151-179	2006
Fukano, Y. Yoshida, T. Yoshimura, H.	Heme oxygenase-1 gene expression in human alveolar epithelial cells (A549) following exposure to whole cigarette smoke on a direct in vitro exposure system	Experimental and Toxicologic Pathology	57	411-418	2006
Hayakawa, K. Hiramatsu, N. Kasai, A. Kitamura, M. Maeda, S. Yao, J.	High Levels of Dioxin-Like Potential in Cigarette Smoke Evidenced by In vitro and In vivo Biosensing	Cancer Res	66(14)	7143-7150	2006
Hur, G.Y. In, K.H. Jung, H.C. Jung, K.H. Kang, E.J. Kang, K.H. Kim, J.H. Lee, S.Y. Lee, S.Y. Shim, J.J. Shin, C. Yoo, S.H.	Peroxisome proliferator-activated receptor- γ inhibits cigarette smoke solution-induced mucin production in human airway epithelial (NCI-H292) cells	Am J Physiol Lung Cell Mol Physiol	291	84-90	2006
Jumarie, C. Morin, A. Porter, A. Prefontaine, D.	In vitro bioactivity of combustion products from 12 tobacco constituents	Food and Chemical Toxicology	44	724-738	2006

Kropotov, A. Krutilina, R. Leutenegger, C. Pleskach, N. Serikov, V.B. Suh, J.H. Tomilin, N.V.	Cigarette Smoke Extract Inhibits Expression of Peroxiredoxin V and Increases Airway Epithelial Permeability	Inhalation Toxicology	18	79-92	2006
Lin, S. Talbot, P. Wu, M. Yu, R.	Cigarette Smoke Toxicants Alter Growth and Survival of Cultured Mammalian Cells	Toxicological Sciences	93(1)	82-95	2006
Mendes, E. Randhawa, K. Wanner, A.	Acute Effect of Cigarette Smoke and Nicotine on Airway Blood Flow and Airflow in Healthy Smokers	Lung	184	363-368	2006
Renne RA, Yoshimura H, Yoshino K, Lulham G, Minamisawa S, Tribukait A, Dietz DD, Lee KM & Westerberg RB	Effects of flavouring and casing ingredients on the toxicity of mainstream cigarette smoke in rats.	Inhalation Toxicology	18	685-706	2006
Aliaga, C. Dennis, P.A. Desai, D. El-Bayoumy, K. Fiala, E.S. Reinhardt, J. Seibert, E. Sodum, R.S. Sohn, O.S. Tsurutani, J. Wang, C.-X.	Induction of preneoplastic lung lesions in guinea pigs by cigarette smoke inhalation and their exacerbation by high dietary levels of vitamins C and E	Carcinogenesis	26(3)	605-612	2005
Arts, J.H.E. Buttner, A. Hausmann, H.-J. Stinn, W.	Cigarette Mainstream Smoke and Gas Phase-Depleted Particulate Phase Enhance Lung Tumorigenicity in the A/J Mouse	SOT Conference, New Orleans, 2005	Poster New Orleans 05	2	2005
Barr, E.B. Belinsky, S.A. Finch, G.L. Gigliotti, A.P. Grimes, M.J. Hahn, F.F. Hobbs, C.H. Hutt, J.A. March, T.H. Mauderly, J.L. Seilkop, S.K. Vuilleminot, B. R.	Life-span inhalation exposure to mainstream cigarette smoke induces lung cancer in B6C3F1 mice through genetic and epigenetic pathways	Carcinogenesis	26(11)	1999-2009	2005

Bartalesi, B. Cavarra, E. Fineschi, S. Lucattelli, M. Lungarella, G. Lunghi, B. Martorana, P.A.	Different lung responses to cigarette smoke in two strains of mice sensitive to oxidants	Eur Respir J	25 (1)	15-22	2005
Bombick, B.R. Doolittle, D.J. Fields, W.R.	Gene Expression in Bronchial Epithelial Cells Following In Vitro Exposure to Cigarette Smoke Condensate	CORESTA 2005	SSPT23	16-Jan	2005
Bottlaender, M. Coulon, C. Dolle, F. Ottaviani, M. Syrota, A. Valette, H.	Acute inhibitor of Cardiac Monoamines Oxidase A after Tobacco Smoke Inhalation: Validation Study of [11C] Befloxatone in Rats Followed by a Positron Emission Tomography Application in Baboons.	The Journal of Pharmacology and Experimental Therapeutics	314(1)	431-436	2005
Boyle, J.O. Dannenberg, A.J. Duffield-Lillico, A.J. Gross, N.D. Morrow, J.D. Moskowitz, C.S. Subbaramaiah, K. Williams, M.K.	Levels of Prostaglandin E Metabolite, the Major Urinary Metabolite of Prostaglandin EZ, Are Increased in Smokers	Clin Cancer Res.	11(16)	6087-6093	2005
Chaudhuri, S.P. John, L. Pillai, B. Sharma, G.	Cigarette smoke extract induces changes in growth and gene expression of <i>Saccharomyces cerevisiae</i>	Biochemical and Biophysical Research Communications	338	1578-1586	2005
Chen, K. Chen, L. Gao, S. Li, W. Rich, C.B. Stone, P. Zhao, Y.	Transcriptional and Posttranscriptional Inhibition of Lysyl Oxidase Expression by Cigarette Smoke Condensate in Cultured Rat Fetal Lung Fibroblasts	Toxicological Sciences	87(1)	197-203	2005

Dawe, D.E. Fattouh, R. Inman, M.D. Jordana, M. Pouladi, M.A. Richards, C.D. Robbins, C.S. Stampfli, M.R. Vujcic, N.	Mainstream Cigarette Smoke Exposure Attenuates Airway Immune Inflammatory Responses to Surrogate and Common Environmental Allergens in Mice, Despite Evidence of Increased Systemic sensitization	Journal of Immunology	175	2834-2842	2005
Ejaz, S. Seok, K.B. Woong, L.C.	Toxicological Effects of Mainstream Whole Smoke Solutions on Embryonic Movements of the Developing Embryo	Drug and Chemical Toxicology	1	14-Jan	2005
Gottschalk, R.W.H. Kleinjans, J.C.S. Moonen, E.J. van Delft, J.H.M. van Herwijnen, M.H. van Leeuwen, D.M.	Differential Gene Expression in Human Peripheral Blood Mononuclear Cells Induced by Cigarette Smoke and Its Constituents	Toxicological Sciences	86(1)	200-210	2005
Hamm, J.T. Misra, M. Morrissey, R.L. Rajendran, N. Yee, S.F.	Focal Proliferative Lesions in A/J Mouse Lung Following 5-Month Exposure to Mainstream Cigarette Smoke	SOT Conference, New Orleans, 2005	PosterNew Orleans 05	2	2005
Hartman, H. Holloway, G. Millet, S. Sheabar, F. Ye, J.	Evaluation of Cigarette Smoke Cytotoxicity Under In Vitro Conditions	SOT Conference, New Orleans, 2005	SOT Poster March 05	1	2005
Hausmann, H.-J. Meurrens, K. Ruf, S. Schleef, R. Schluter, K.-D. von Holt, K.	Influence of Sub- Chronic Cigarette Smoke Exposure on the Progression of Myocardial Hypertrophy in Spontaneously Hypertensive Rats (SHR)	SOT Conference, New Orleans, 2005	Poster NewOrlean s 05	2	2005
Heuer, L.S. Kaul, R. Pinkerton, K.E. Thao, N.N. Wenman, W.M. Wiedeman, J.A.	Tobacco smoke induces a persistent, but recoverable state in Chlamydia pneumoniae infection of human endothelial cells	Microbial Pathogenesis	39	197-204	2005

Hirano, Y. Ishii, H. Kobayashi, M. Kubo, S. Masunaga, Y. Shimizu, Y. Takahashi, K.	Cytokine and chemokine expression in cigarette smoke-induced lung injury in guinea pigs	Eur Respir J	26	993-1001	2005
Hoffmann, H. Hogel, J. Isner, C. Speit, G.	Genetic polymorphisms and the effect of cigarette smoking in the comet assay	Mutagenesis	20(3)	359-364	2005
Witschi, H.	Carcinogenic Activity of Cigarette Smoke Gas Phase and Its Modulation by Beta-Carotene and N-Acetylcysteine	Toxicological Sciences	84	81-87	2005
Zhong, K.	Effect of Smoking on the Volatile SulfuretedCompounds (VSCs)inOral Cavity	CORESTA 2005	SSPT37	16-Jan	2005
Ahmad, A. Allen, R.H. Panayiotidis, M.I. Stabler, S.P. White, C.W.	Cigarette smoke extract increases S-adenosylmethionine and cystathionine in human lung epithelial-like (A549) cells	Chemico-Biological Interactions	147	87-97	2004
Aufderheide, M. Knebel, J.W. Ritter, D.	Comparative Assessment of Toxicities of Mainstream Smoke from Commercial Cigarettes	Inhalation Toxicology	16	691-700	2004
Aufderheide, M. Mohr, U.	The use of the modified CULTEX system for the direct exposure of bacteria to mainstream cigarette smoke	Coresta Congress, Kyoto, Japan		08-Jan	2004
Ayres, P.H. Curtin, G.M. Higuchi, M.A. Mosberg, A.T. Swauger, J.E.	Lung Tumorigenicity in A/J and rasH2 Transgenic Mice Following Mainstream Tobacco Smoke Inhalation	Toxicological Sciences	81	26-34	2004
Ayres, P.H. Higuchi, M.A. Sagartz, J.W. Shreve, W.K.	Comparative Subchronic Inhalation Study of Smoke From the 1R4F and 2R4F Reference Cigarettes	Inhalation Toxicology	16	20-Jan	2004
Baker RR, Massey ED & Smith G	An overview of the effects of tobacco ingredients on smoke chemistry and toxicity	Food and Chemical Toxicology	42	53-83	2004

Barakat, A.I. Douglas, G.R. Mariano, N.F. Soghomonians, A. Thirkill, T.L.	Effect of Aqueous Tobacco Smoke Extract and Shear Stress on PECAM-1 Expression and Cell Motility in Human Uterine Endothelial Cells	Toxicological Sciences	81	408-418	2004
Barankin, B. Bird, G. Freiman, A. Lauzon, G.J. Metelitsa, A.I.	Cutaneous Effects of Smoking	Journal of Cutaneous Medicine and Surgery	8(6)	415-413	2004
Barr, E.B. Bechtold, W. E. Belinsky, S.A. Finch, G.L. Gigliotti, A.P. Hahn, F.F. Hobbs, C.A. March, T.H. Mauderly, J.L. Seilkop, S.K.	Chronic Inhalation Exposure to Mainstream Cigarette Smoke Increases Lung and Nasal Tumor Incidence in Rats	Toxicological Sciences	81	280-292	2004
Bosio, A. Gebel, S. Gerstmayer, B. Haussmann, H.-J. Muller, T. Van Miert, E.	Gene expression profiling in respiratory tissues from rats exposed to mainstream cigarette smoke	Carcinogenesis	25 no 2	169-178	2004
Carchman, R.A. Gaworski, C.L. Meisgen, T.J. Podraza, K.F. Reininghaus, W. Roemer, E. Rustemeier, K. Stabbert, R. Veltel, D.J.	Chemical composition, cytotoxicity and mutagenicity of smoke from US commercial and reference cigarettes smoked under two sets of machine smoking conditions	Toxicology	195	31-52	2004
Chaung, Y.H. Chen, H.W. Chien, M.L. Lii, C.K. Wang, T.S.	Extracts from cigarette smoke induce DNA damage and cell adhesion molecule expression through different pathways	Chem.-Biol. Interactions	150	233-241	2004
Chen, D. Hayata, I. Minamihisamatsu, M. Morishima, H. Sugahara, T. Wang, C. Wei, L. Yuan, Y. Zhang, W.	Effect of Smoking on Chromosomes Compared with That of Radiation in the Residents of a High-Background Radiation Area in China	J. Radiat. Res.	45 (3)	441-446	2004

Curtin, G.M. Hanausek, M. Mosberg, A.T. Slaga, T.J. Walaszek, Z.	Short-Term In Vitro and In Vivo Analyses for Assessing the Tumor-Promoting Potentials of Cigarette Smoke Condensates	Toxicological Sciences	81	14-25	2004
DeMarini, D.M.	Genotoxicity of tobacco smoke and tobacco smoke condensate: a review	Mutation Research	567	447-474	2004
Dolas, S.S. Maru, G. B. Pakhale, S.S. Thapliyal, R.	Evaluation of DNA Damage in Mice Topically Exposed to Total Particulate Matter from Mainstream and Sidestream Smoke from Cigarettes and Bidis	Mutagenesis	19	413-421	2004
Fuciarelli, A.F. Lee, K.-M. McKinney, W.J. Obot, C. J. Renne, R.A.	Characterization of Mainstream Cigarette Smoke-Induced Biomarker Responses in ICR and C576I/6 Mice	Inhalation Toxicology	16	701-719	2004
Garg, R. Hansch, C. Martin, P. Perfetti, T.A. Smith, C.J.	Percutaneous penetration enhancers in cigarette mainstream smoke	Food and Chemical Toxicology	42	15-Sep	2004
Geerlings, M. Hylkema, M.N. Kerstjens, H.A.M. Klok, P. A. Luinge, M.A. Melgert, B.N. Postma, D.S. Timens, W. van der Strate, B.W.A.	Short-Term Smoke Exposure Attenuates Ovalbumin-Induced Airway Inflammation In Allergic Mice	Am J Respir Cell Mol Biol	30	880-885	2004
Goldstraw, P. Pastorino, U. Ratcliffe, C. Tetley, T.D. Vanden Bon, E.J. Witherden, I.R.	Primary Human Alveolar Type II Epithelial Cell Chemokine Release Effects of Cigarette Smoke and Neutrophil Elastase	Am J Respir Cell Mol Biol	30	500-509	2004
Hayes, J.R. Meckley, D.R. Mosberg, A.T. Swauger, J.E. Van Kampen, K.R.	A responsive, sensitive, and reproducible dermal tumor promotion assay for the comparative evaluation of cigarette smoke condensates	Regulatory Toxicology and Pharmacology	39	135-149	2004

Hirose, M. Lee, I.S. Mori, Y. Nishikawa, A. Tanaka, T.	Cigarette Smoking, Metabolic Activation and Carcinogenesis	Current Drug Metabolism	5 (5)	363-373	2004
Intorp, M. Roper, W. Wieczorek, R.	HOFFMANN ANALYTES AND CIGARETTE SMOKE IN VITRO TOXICITY REVISITED - HOW DO THE DATA COMPARE?	Coresta Congress, Kyoto, Japan		08-Jan	2004
Samet, J.M.	Adverse effects of smoke exposure on the upper airway	Tob. Control	13 sup.1	57-60	2004
Abbott, B.D. Rogers, J.M.	Screening for Developmental Toxicity of Tobacco Smoke Constituents	Toxicological Sciences	75	227-228	2003
Aggarwal, B.B. Gairola, C.G. Potdar, P. Shishodia, S.	Curcumin (diferuloylmethane) down-regulates cigarette smoke- induced NF-kB activation through inhibition of Ikb α kinase in human lung epithelial cells:correlation with suppression of COX-2, ...	Carcinogenesis	24(7)	1269-1279	2003
Andreoli, C. Gigante, D. Nunziata, A.	A review of in vitro methods to assess the biological activity of tobacco smoke with the aim of reducing the toxicity of smoke	Toxicology in Vitro	17	587-594	2003
Aoshiba, K. Koinuma, M. Nagai, A. Yokohori, N.	IMMUNOHISTOCHEMI CAL EVALUATION OF OXIDATIVE STRESS IN MURINE LUNGS AFTER CIGARETTE SMOKE EXPOSURE	Inhalation Toxicology	15	1029-1038	2003
Arey, J. Iv, M. Riveles, K. Talbot, P.	Pyridines in cigarette smoke inhibit hamster oviductal functioning in picomolar doses	Reproductive Toxicology	17	191-202	2003

Aufderheide, M. Knebel, J.W. Ritter, D.	EXPOSURE OF HUMAN LUNG CELLS TO INHALABLE SUBSTANCES: A NOVEL TEST STRATEGY INVOLVING CLEAN AIR EXPOSURE PERIODS...	Inhalation Toxicology	15	67-84	2003
Ayres, P.H. Kinsler, S. Mosberg, A.T. Pence, D.H. Shreve, W.K.	RAT SUBCHRONIC INHALATION STUDY OF SMOKE FROM CIGARETTES CONTAINING FLUE- CURED TOBACCO CURED EITHER BY DIRECT-FIRED OR HEAT-EXCHANGER CURING PROCESSES	Inhalation Toxicology	15	819-854	2003
Bagnasco, M. Balansky, R.M. Bennicelli, C. Camoirano, A. Cartiglia, C. D'Agostini, F. De Flora, S. Izzotti, A. Longobardi, M.G. Lubet, R.A. Tampa, E.	Modulation of cigarette smoke-related end- points in mutagenesis and carcinogenesis	Mutation Research	523-524	237-252	2003
Balansky, R.M. Bennicelli, C. Camoirano, A. D'Agostini, F. De Flora, S. Izzotti, A. Lubet, R.A. Wang, Y. You, M. Zhang, Z.Q.	Molecular Alterations and Lung Tumors in 53 Mutant Mice Exposed to Cigarette Smoke	Cancer Res	63	793-800	2003
Barnes, P.J. Buttery, L.D.K. Habib, M. Hoyt, J.C. Polak, J.M. Robbins, R.A. Springall, D.R.	CIGARETTE SMOKE DECREASES INDUCIBLE NITRIC OXIDE SYNTHASE IN LUNG EPITHELIAL CELLS	Experimental Lung Research	29	12	2003

Bosio, A. Bulles, H. Friedrichs, D. Gebel, S. Gerstmayer, B. Koeser, A. Schlage, W.K.	Gene Expression Profiling of Two 3T3 Cell lines Exposed to Cigarette Smoke Fractions Reveals Differences in the Induction of Xenobiotic-Metabolizing Enzymes	9th International Inhalation Symposium (June 11-14) Hannover Germany	June	1	2003
Budhiraja, R. Chalkley, R. Cooray, S. Hassour, P.M. Kayyali, U.S. Lanzillo, J.J. Pennella, C.M.	Upregulation of xanthine oxidase by tobacco smoke condensate in pulmonary endothelial cells	Toxicology and Applied Pharmacology	188	59-68	2003
Dybing, E. Fowles, J.	Application of toxicological risk assessment principles to the chemical constituents of cigarette smoke	Tob. Control	12	424-430	2003
Fowler, C.J. Sandberg, M. Tiger, G.	Effects of Water-Soluble Cigarette Smoke Extracts Upon The Release of Beta-Hexosaminidase from RBL-2H3 Basophilic Leukaemia Cells In Response To Substance P Compound 48-80 Concanavalin A and Antigen	Inflammation Research	52	461-469	2003
Green, C.R. Rodgman, A.	Toxic Chemicals in Cigarette Mainstream Smoke – Hazard and Hoopla (part1)	Beiträge zur Tabakforschung	20 no 8 (part1)	481-545	2003
Green, C.R. Rodgman, A.	Toxic Chemicals in Cigarette Mainstream Smoke - Hazard and Hoopla (part 2)	Beiträge zur Tabakforschung	20 no 8 (part 2)	481-545	2003
Green, C.R. Rodgman, A.	Toxic Chemicals in Cigarette Mainstream Smoke - Hazard and Hoopla (part 3)	Beiträge zur Tabakforschung	20 no 8 (part 3)	481-545	2003

Ishihara, A. Kita, T. Nakashima, T. Nakatani, T.	EFFECTS OF EXPOSURE TO CIGARETTE SMOKE AT DIFFERENT DOSE LEVELS ON EXTENSOR DIGITORUM LONGUS MUSCLE FIBRES IN WISTAR-KYOTO AND SPONTANEOUSLY HYPERTENSIVE RATS	Clinical and Experimental Pharmacology and Physiology	30	671-677	2003
Ortug, C.	Scanning electron microscopic findings in respiratory nasal mucosa following cigarette smoke exposure in rats	Annals of Anatomy	185	207-210	2003
Altuntas, I. Dane, S. Gumustekin, K.	EFFECTS OF CIGARETTE SMOKING ON LIPID PEROXIDATION	Journal of Basic and Clinical Physiology and Pharmacology	13(1)	69-73	2002
Bosio, A. Gebel, S. Hausmann, H.-J. Janssen, U. Knorr, C. Muller, T.	Kinetics of gene expression profiling in Swiss 3T3 cells exposed to aqueous extracts of cigarette smoke	Carcinogenesis	23 no 5	741-748	2002
Carmines EL	Evaluation of the potential effects of ingredients added to cigarettes. Part 1: Cigarette design, testing approach and review of results.	Food and Chemical Toxicology	40	77-91	2002
Coggins, C.R.E.	A MINIREVIEW OF CHRONIC ANIMAL INHALATION STUDIES WITH MAINSTREAM CIGARETTE SMOKE	Inhalation Toxicology	14	991-1002	2002
Degrell, P. Mazak, I. Molnar, G.A. Nagy, J. Vas, T. Wagner, L. Wagner, Z. Wittmann, I.	Cigarette Smoke and Its Formaldehyde Component Inhibit Bradykinin-Induced Calcium Increase in Pig Aortic Endothelial Cells	Endothelium	9	103-108	2002
Garg, R. Hansch, C. Morton, M.J. Perfetti, T.A. Rodgman, A. Selassie, C.D. Smith, C.J.	The Relative Toxicity of Substituted Phenols Reported in Cigarette Mainstream Smoke	Toxicological Sciences	69	265-278	2002

Haginaka, J. Kagota, S. Kunitomo, M. Yamaguchi, Y.	Participation of peroxynitrite in oxidative modification of LDL by aqueous extracts of cigarette smoke	FEBS Letters	512	218-222	2002
Krause, G. Scherer, G. Wolz, L.	The Comet Assay with MCL-5 Cells as an Indicator of Genotoxic Treatment with Chemicals and Cigarette Smoke Condensate	ATLA	30	331-339	2002
Paschke T, Scherer G & Heller W	Effects of ingredients on cigarette smoke composition and biological activity: A literature overview.	Beitrage zur Tabakforschung International	20	107-247	2002
RODGMAN A.	Some studies of the effects of additives on cigarette mainstream smoke properties. II. Casing materials and humectants.	Int.Contributions to Tobacco Research	20-4	279-300.,	2002
Rustemeier K Stabbert R Hausmann H J Roemer E L carmines E L	Evaluation of the potential effects if ingredients added to cigarettes. Part 2: Chemical composition of mainstream smoke.	Food and Chemical Toxicology	40	93-104	2002
Aoki, N. Hasumura, Y. Hioki, Y. Homori, M. Ishikawa, K. Kawaro, K. Maki, A. Yanagisawa, A. Yasumura, Y. Yoshiro, H.	Acute effects of cigarette smoking on platelet-dependent thrombin generation	European Heart Journal	22	56-61	2001
Coggins, C.R.E.	A Review of Chronic Inhalation Studies with Mainstream Cigarette Smoke, in Hamsters, Dogs, and Nonhuman Primates	Toxicologic Pathology	29	550-557	2001
Avalos, J.T. Bombick, B.R. Borgerding, M.F. Chepiga, T.A. Doolittle, D.J. Morton, M.J. Murphey, P.A. Swager, J.E.	A comparison of the mainstream smoke chemistry and mutagenicity.,of a representative sample of the US cigarette market with two Kentucky reference cigarettes (KI R4F and KI R5F)	Food and Chemical Toxicology	38	949-962	2000

Hansch, C. Smith, C.J.	The Relative Toxicity of Compounds in Mainstream Cigarette Smoke Condensate	Food and Chemical Toxicology	38	637-646	2000
Carchman, R.A. Gomm, W.A. Meisgen, T.J. Roemer, E. Tewes, F.J.	CIGARETTE PARAMETERS THAT INFLUENCE THE MUTA-GENICITY OF MAINSTREAM SMOKE CONDENSATE.	Toxicological Sciences	48	125	1999
Gaworski CL, Heck JD, Bennett MB & Wenk ML	Toxicologic evaluation of flavour ingredients added to cigarette tobacco: skin painting bioassay of cigarette smoke condensate in SENCAR mice.	Toxicology	139	1-17	1999
Aufderheide, M. Jarck, P. Knebel, J.W. Koch, W. Lodding, H. Massey, E. Pohlmann, G. Windt, H.	Micronucleus induction in V79 cells after direct exposure to whole cigarette smoke	Mutagenesis	13(2)	145-149	1998
Gaworski CL, Dozier HM, Heck JD, Gerhart JM, Rajendran N, David RM, Brennecke LH & Morrissey R	Toxicologic evaluation of flavour ingredients added to cigarette tobacco: 13-week inhalation exposures in rats.	Inhalation Toxicology	10	357-381	1998
Ashida, K. Itoh, T. Kamisaki, Y. Kishimoto, Y. Nakamoto, K. Wada, K.	SUBSTANCES IN THE AQUEOUS FRACTION OF CIGARETTE SMOKE INHIBIT LIPID PEROXIDATION IN SYNAPTOSOMES OF RAT CEREBRAL CORTEX	Biochemistry and Molecular Biology International	42(1)	10-Jan	1997
Ayres, P.H. Bombick, D.W. Doolittle, D.J.	CYTOTOXICITY ASSESSMENT OF WHOLE SMOKE AND VAPOR PHASE OF MAINSTREAM AND SIDESTREAM CIGARETTE SMOKE FROM THREE KENTUCKY REFERENCE CIGARETTES	Toxicology Methods	7	177-190	1997
Ohshima, H. Yoshie, Y.	Synergistic induction of DNA strand breakage by cigarette tar and nitric oxide	Carcinogenesis	18(7)	1359-1363	1997

Kalra, V.K. Rattan, V. Shen, Y. Sultana, C.	Cigarette smoke condensate-induced adhesion molecule expression and transendothelial migration of monocytes	American Journal of Physiology	270	1624-1633	1996
Doolittle, D.J. Fulp, C.W. Lee, C.K. Payne, V.M. Rees, D.C. Steele, R.H.	A comparison of the mutagenicity of mainstream cigarette smoke condensates from a representative sample of the U.S. cigarette market with a Kentucky reference cigarette (KIR4F)	Mutation Research	342	179-190	1995
Eiro, H.	EFFECTS OF CIGARETTE TOBACCO TAR BY GASTRIC ADMINISTRATION ON MURINE PULMONARY TUMORIGENESIS	Journal of Toxicologic Pathology	8	377-384	1995
Burns, A.R. Dunn, L.A. Hogg, J.C. Hosford, S.P. Walker, D.C.	Respiratory epithelial permeability after cigarette smoke exposure in guinea pigs	J Appl Physiol	66(5)	2109-2116	1989
Gonzalez-Rothi, R.J. Harris, J.O.	EFFECTS OF LOW-YIELD-CIGARETTE SMOKE INHALATION ON RAT LUNG MACROPHAGES	Journal of Toxicology and Environmental Health	17	221-228	1986
Hudgens, R.W. Sonnenfeld, G.	Effect of Sidestream and Mainstream Smoke Exposure on in Vitro Interferon - alpha/beta Production by L-929 Cells	Cancer Res	46	2779-2783	1986
Schmahl, D. Zeller, W.J.	Relevance of Gas and Particulate Phases of Tobacco Smoke for Lung Cancer Formation	Cancer Detection and Prevention	9	91-97	1986
Gairola, C.	Genetic effects of fresh cigarette smoke in <i>Saccharomyces cerevisiae</i>	Mutation Research	102	123-136	1982
Boyd, C.H. Dansie, D.R. Henry, C.J. Kouri, R.E. Rasmussen, R.E.	DNA Replication and Unscheduled DNA Synthesis in Lungs of Mice Exposed to Cigarette Smoke	Cancer Res	41	2583-2588	1981

Coggins, C.R.E. Fouillet, X.L.M. Lam, R. Morgan, K.T.	CIGARETTE SMOKE INDUCED PATHOLOGY OF THE RAT RESPIRATORY TRACT: A COMPARISON OF THE EFFECTS OF THE PARTICULATE AND VAPOUR PHASES	Toxicology	16	83-101	1980
Lam, R.	TRANSIENT EPITHELIAL LOSS IN RAT LARYNX AFTER ACUTE EXPOSURE TO TOBACCO SMOKE	Toxicology Letters	6	327-335	1980
Bernfeld, P. Homburger, F. Pai, K.J. Soto, E.	Cigarette Smoke Inhalation Studies in Inbred Syrian Golden Hamsters	J Natl Cancer Inst (JNCI)	63(3)	675-689	1979
Akin, F. J. Benner, J.F.	Induction of Aryl Hydrocarbon Hydroxylase in Rodent Lung by Cigarette Smoke : A Potential Short-Term Bioassay	Toxicology and Applied Pharmacology	36	331-337	1976
Ames, B.N. Kier, L. D. Yamasaki, E.	Detection of Mutagenic Activity in Cigarette Smoke Condensates	Proc. Natn. Acad. Sci. USA	71 no 10	4159-4163	1974
Leuchtenberger, C. Leuchtenberger, R. Zbinden, I.	Gas Vapour Phase Constituents and SH Reactivity of Cigarette Smoke Influence Lung Cultures	Nature	247	565-567	1974
Chevalier, H.-J. Dontenwill, W. Harke, H.-P. Lafrenz, U. Reckzeh, G. Schneider, B.	Investigations on the Effects of Chronic Cigarette-Smoke Inhalation in Syrian Golden Hamsters	J Natl Cancer Inst (JNCI)	51	1781-1832	1973
Dontenwill, W. Elmenhorst, H. Harke, H.-P. Misfeld, J. Reckzeh, G. Timm, J. Weber, K.H.	Experimentelle Untersuchungen über die tumorerzeugende Wirkung von Zigarettenrauch- Kondensaten an der Mäusehaut	Z. Krebsforsch.	73	285-304	1970

Leuchtenberger, C. Leuchtenberger, R.	Cytologic and Cytochemical Effects on Primary Mouse Kidney Tissue and Lung Organ Cultures after Exposure to Whole Fresh Smoke	Cancer Res	29	862-872	1969
Benedict, R. C. Stedman, R.L.	Complexity of Enzymatic Inhibition by Cigarette Smoke	Experientia	24 no 12	1205-1206	1968



Polycyclic aromatic hydrocarbons (Benzo[a]pyrene) Toxicological Overview

Key Points

Kinetics and metabolism

- PAHs are absorbed by all routes of exposure
- PAHs are distributed widely throughout the body, with fatty tissues tending to show higher amounts
- metabolites of PAH's are generally excreted as conjugates of GSH, glucuronic acid or sulphate in the urine, faeces and via biliary excretion

Health effects of acute exposure

- few studies were identified that reported the effects of BaP alone in humans following acute inhalation, ingestion or dermal exposure

Health effects of chronic exposure

- chronic exposure to mixtures of PAHs in air have resulted in a range of respiratory effects, ischemic heart disease, chronic dermatitis, depressed immune system and cancer of the skin and lungs
- BaP amongst other PAHs is able to form DNA adducts which are likely key to their mutagenic potential
- complex mixtures of PAHs which include BaP are considered to be carcinogenic to humans

Summary of Health Effects

PAHs typically occur in complex mixtures and not as individual compounds, BaP is considered to be one of the most toxic PAHs and has been extensively studied. For the general public, the main route of exposure to poly aromatic hydrocarbons (PAHs) is ingestion of food. For smokers, the contribution of smoking to total PAH exposure will be similar to that of food. Inhalation and skin absorption are the main routes of occupational exposure.

PAHs are rapidly distributed throughout the body following all routes of exposure; they may be detected in most tissues minutes to hours following exposure. Benzo[a]Pyrene (BaP) is metabolised by cytochrome P450 enzymes and epoxide hydrolase resulting in a number of metabolites being formed. These metabolites include the reactive epoxide BaP 7,8 diol-9,10-epoxide (BPDE), which is believed to play a role in the carcinogenicity of BaP. Following metabolism PAHs are excreted in the urine and/or faeces depending on their molecular weight.

No data on the acute effects of BaP in humans were identified and few studies were reported in animals. Following acute oral exposure of rats to BaP, effects on the liver were observed.

Following chronic exposure to PAHs in an occupational setting a decrease in lung function was reported, as well as chest pain, respiratory irritation, cough, dermatitis and depressed immune system; although in most cases it was not possible to evaluate the contribution of BaP to such effects. Few adverse effects were observed in rats or hamsters exposed to BaP via inhalation. Following ingestion, myelotoxicity was observed in poor affinity aryl hydrocarbon-receptor (AhR) mice but not in high affinity mice. Hepatotoxicity was also reported.

BaP can cross the placenta; BaP-DNA adducts have been found in foetal cord blood and also in the sperm. Presence of these adducts has been associated with reduced sperm count in men and decreased motor development in infants. BaP was found to cause adverse developmental and reproductive effects in mice and rats. Dietary administration during gestation reduced fertility and fetal abnormalities whereas administration by gavage caused an increase in fetal death and decreased fertility.

Biomarkers of exposure to BaP are seen concurrently with biomarkers of genotoxic effect. The International Agency for Research on Cancer (IARC) has stated that BaP likely contributes to the genotoxic action of complex PAH mixtures that individuals may be exposed to occupationally.

IARC has classified BaP as carcinogenic to humans (Group 1). No epidemiological studies of exposure to BaP alone in man can be identified; however sufficient data exists for exposure to complex PAH mixtures containing BaP.

Introduction

PAHs are a large group of compounds consisting of hydrocarbons containing two or more benzene rings fused together or to other hydrocarbon rings. They are formed during the combustion of carbonaceous material at high temperatures [1]. PAHs typically occur in complex mixtures and not as individual compounds, the process generating the mixture may define its composition. BaP is considered to be one of the most toxic PAHs and has therefore been extensively studied; hence the majority of the data in this document refers to BaP. Unless otherwise specified, "PAHs" will refer to a mixture of PAH compounds (which may include BaP).

Kinetics and Metabolism

PAHs are lipophilic compounds that are readily absorbed from the lungs following inhalation, the gastrointestinal (GI) tract following ingestion and the skin following dermal exposure [2].

PAHs may adsorb onto particulate matter in air. In humans, it was reported that BaP measured in the lungs following inhalation of soot particles was much lower than expected. This may be due to the ability of the pulmonary epithelial cells to metabolise BaP thereby facilitating its absorption and clearance from the lungs [3]. Occupational studies have inferred that inhaled PAHs are absorbed, as urinary metabolites were present in workers exposed to PAHs [2]. The absorption of BaP following inhalation is highly dependent on the type of particles onto which it is adsorbed and the site of deposition in the respiratory tract [1]. Pulmonary absorption often occurs in parallel with mucociliary clearance, by which PAHs that are absorbed onto inhaled particulates are cleared out of the pulmonary tree and subsequently swallowed [3, 4].

PAHs are well absorbed in the GI tract by passive diffusion. The extent of absorption may vary depending of the bioavailability of particles, diet and PAH size (highest for smaller molecule PAHs) [1]. In rats, approximately 40% of absorption occurred through the GI tract following administration of 0.5 µg/kg BaP for 90 minutes directly into the duodenum; 38-58% absorption occurred following administration of BaP given by gavage or in the diet [2].

BaP is efficiently absorbed through the skin of animals [1]. Extensive skin absorption has been demonstrated in mice, as almost all of an applied dose of BaP appeared in the faeces following application to the skin [4]. Similarly, rapid absorption was demonstrated in rats, monkeys and guinea pigs [2].

PAHs are rapidly distributed throughout the body by all routes of exposure; they may be detected in most tissues minutes to hours following exposure [1]. Fatty tissues generally contain more PAHs than other tissues [1, 5]. However, PAHs do not accumulate in the body [1].

BaP can readily cross the placenta following oral, inhalation or dermal exposure. One study reported that when pregnant rats were exposed to BaP via inhalation, an increase in BaP and metabolites was measured in both maternal and fetal blood and tissues. Similarly, BaP

was measured in the fetus when rats were given oral BaP on day 21 of pregnancy [2]. BaP has been detected in human breast milk, particularly in smokers, however animal studies suggest that it does not accumulate in the offspring [6].

Cytochrome P450 enzymes (CYPs) and epoxide hydrolase are the main enzymes involved in PAH metabolism. Many studies have investigated the metabolism of PAHs in tissues and cells following ingestion of food containing PAHs, or inhalation or ingestion of environmental PAHs. The liver has the greatest capacity to metabolise PAHs followed by the lungs, intestinal mucosa, skin and kidneys. BaP stimulates its own metabolism by inducing CYP enzymes via the activation of the AhR. PAHs can also inhibit CYP enzymes [1].

PAHs are initially metabolised to several epoxides by CYP enzymes. The epoxides may spontaneously rearrange to phenols or are converted to dihydrodiols. PAHs are also metabolised to a number of quinones by CYPs. The dihydrodiols are further metabolised by CYPs to 4 optically active isoforms of dihydrodiol epoxides, notable amongst these is the anti-BaP 7,8 diol-9,10-epoxide (anti-BPDE). The stereoisomer (+) anti-BPDE is considered to be the most tumorigenic and predominant metabolite of BaP that forms DNA adducts in mammalian tissues. In addition, PAHs and their reactive metabolites undergo conjugation with sulphate, glutathione (GSH) and glucuronic acid [1, 7].

The route of exposure may influence the toxicity of PAHs. Following oral exposure the compound undergoes first-pass metabolism which reduces the levels of PAHs and metabolites that reach systemic circulation. Inhalation or dermal exposure may result in higher levels of PAHs reaching peripheral tissues as the compounds may bypass the first-pass effect of the liver [1, 2]. Genetic polymorphisms may affect the capacity of individuals to metabolise PAHs [1].

Studies suggest that metabolites of PAH's are generally excreted as conjugates of GSH, glucuronic acid or sulphate in the urine, faeces and via biliary excretion [7]. High molecular weight PAHs and their metabolites are mainly excreted via the faeces [1].

Sources and Route of Human Exposure

The major route of exposure to PAHs for non-smokers in the general population is food, with a minor contribution from inhalation of ambient air. For smokers, the contribution of smoking to total exposure is likely to be similar to that of food [8].

Food may become contaminated with PAHs from environmental sources, industrial preparation or during home cooking [8]. Various foods such as vegetables, meat and fish have been shown to contain PAHs, but they are largely formed due to cooking at high temperatures such as charbroiling, grilling and frying. Smoked and barbequed food are particularly important sources of exposure, although the largest contributors to PAH intake are "cereals and cereal products" and "vegetable fats and oils" [3, 4, 9-11]. The maximum estimated daily intake of BaP for a 70 kg person is 6-8 ng/kg [8, 11]. After evaluating a recent food survey, the Food Standards Agency (FSA) concluded that PAHs were typically

found in low levels in food and that consumers do not need to change their eating habits [12].

In the past, metal production and agriculture were responsible for the majority of total BaP emissions in the UK. Likely effected by the Environmental Protection Act 1990 and the ban on burning agricultural stubble, total emissions in 2011 had reduced to roughly 1/20th (now 3 tonnes) of those in 1991. Natural sources in 2011 accounted from 47.2% of the total emissions, while residential and commercial sources contributed the greatest portion of the anthropogenic emissions at 76% [13].

Annual mean air concentrations of BaP in the UK are generally below the EU target value of 1 ng/m³, with the vast majority of monitoring sites showing levels below the UK air quality objective of 0.25 ng/m³. Locations with point sources (e.g. industrial instillations and domestic solid fuel burning) are an exception to this. The main sources of BaP in the UK are domestic coal and wood burning, outdoor fires and industrial processes [14, 15].

Indoor air may be contaminated with PAH's by infiltration of outdoor air or from indoor emissions, which include smoking, cooking, and heating with fuel stoves and fireplaces and to a lesser extent from incense and candle burning. Levels of BaP within the home appear to vary seasonally, with the highest concentrations found in winter. BaP levels in European homes were found to be between 0.01 to 0.65 ng/m³ [1].

Mainstream tobacco smoke contains high concentrations of PAHs, levels in the range of 1-1.6 µg per cigarette have been measured and as such this represents a major source of exposure for smokers. BaP levels in sidestream smoke have been reported to range from 52-95 ng per cigarette, more than three times higher than that seen in mainstream smoke [5]. In a smoker's home more than 87% of total PAH's in air may be introduced by cigarette smoke; in a room heavily polluted with cigarette smoke, BaP levels may be as high as 22 ng/m³ [1].

PAHs are commonly detected in surface waters, due to urban runoff and industrial activities [3]. Contamination of drinking water with PAHs is usually associated with coal tar linings of distribution pipes. However, drinking water contributes only a minor amount to the total intake of PAHs [4, 16].

PAHs are found in the majority of surface soils due to atmospheric deposition or urban runoff. Soils near industrial sources such as coal coking often contain high concentrations of PAHs [3, 9]. BaP in English soils comprises approximately 5-7% by weight of the total PAH content [17]. The British Geological Survey defined the normal background concentrations for BaP in England and Wales to be 3.6 mg/kg in urban areas and 0.5 mg/kg in all other areas [17, 18].

Occupational exposure is largely through inhalation and skin absorption. The greatest levels of occupational exposure to BaP are in aluminium production (up to 100 µg/m³), with lesser exposure in roofing and paving (10-20 µg/m³) and lesser still in coal processing, wood impregnation, chimney sweeping and in power plants (at or below 1 µg/m³) [5].

Health Effects of Acute/Single Exposure

Human data

Inhalation

No studies were identified that reported the effects of BaP in humans following acute inhalation exposure.

Ingestion

Data on acute oral toxicity of BaP in humans are not available.

Dermal/ocular exposure

No studies were identified that reported effects of BaP in humans following acute dermal exposure.

Animal and in-vitro data

Inhalation

No studies were identified that reported effects of BaP in animals following acute inhalation exposure.

Ingestion

Exposure of rats (intragastric administration) to 100 mg/kg bw/day BaP for four days increased relative liver weight by 27% and induced aldehyde dehydrogenase. Limited evidence suggested that acute ingestion of BaP (50-150 mg/kg bw/day for 4 days) does not cause adverse GI effects in rats, although enzyme activity was altered. It was suggested that more serious effects may occur at higher concentrations [2].

Dermal/ocular exposure

BaP applied dermally caused allergen specific contact hypersensitivity reactions in mice after acute applications of 120 µg. A dose dependent contact hypersensitivity response to dermal application of BaP has been observed in guinea pigs; two applications of 0.001% BaP over 2-3 weeks gave slight hypersensitivity while a 1% dose gave a more severe response [2].

Acute topical application of BaP (concentration and duration of exposure not stated) to the backs of shaved mice suppressed sebaceous glands, although it was not possible to determine if such effects were due to the solvent or BaP, as a control group was not used [2].

Health Effects of Chronic/Repeated Exposure

Human data

Inhalation

A large number of epidemiological studies have been carried out considering a variety of occupations in which the workforce is chronically exposed to a mixture of PAHs [4]. These studies have demonstrated that such exposures result in symptoms including respiratory distress, decreased ventilatory function, chest pain, chest and throat irritation, cough, haematemesis, chronic dermatitis, depressed immune system and cancer of the skin and lungs [2, 4]. It is not possible to determine with any certainty the contribution of individual PAHs to these effects [4].

One study investigated the respiratory effects of inhaled BaP in employees working in various areas of a rubber factory. The authors reported a decrease in ventilatory function following prolonged exposure, as assessed by duration of employment, the greatest effects being observed in workers that had the highest exposure to particulate matter and BaP. No attempt was made to identify other possible chemical exposures or to separate effects due to BaP or particulates [2].

Ischemic heart disease was observed to increase in a dose dependent manner in asphalt workers exposed to BaP. Mean exposure for the cohort was 273 ng/m³ and exposure at or above this level was associated with a 1.64 fold greater risk of ischemic heart disease mortality compared to those exposed to below 68 ng/m³ [1]

Ingestion

Data on chronic oral toxicity of BaP in humans are not available.

Dermal/ocular exposure

Few data are available pertaining to BaP alone.

Regressive verrucae (warts) were reported in humans following up to 120 applications of 1% BaP over a four month period [2].

Genotoxicity

The formation of DNA adducts is believed to be a key event in the mutagenicity and carcinogenicity of PAH's; adducts may lead to misrepair and result in mutations [1]. Anti-BPDE has been demonstrated to form DNA adducts in man and as such acts as a biomarker for exposure to BaP. Molecular epidemiological studies in individuals exposed to complex mixtures of PAHs have shown that BaP adducts are seen concomitantly with biomarkers of genotoxic effect. The observed effects include chromosomal aberrations, sister chromatid exchange, DNA damage and formation of 8-oxo-deoxyguanosine. These same markers of exposure and effect are also observed in experimental animals, with association. IARC considers that BaP contributes to the genotoxic effects seen in complex

PAH mixtures; with the anti-BPDE-DNA adduct being the most mechanistically relevant adduct [19].

Smoking and diet (major sources of BaP) have been highly correlated with levels of the anti-BPDE-DNA adduct. The metabolite responsible for these adducts is seen to cause a unique array of mutations in the TP53 tumour suppressor gene, in cancers associated with smoking [19].

Evidence suggests that inhalation exposure to BaP at levels over 1 ng/m³ is predictive of greater genomic frequency of translocation, micronuclei and DNA fragmentation [1].

Carcinogenicity

No epidemiological studies on exposure to BaP alone are available for evaluation [19]. There is however extensive literature on the epidemiology of workforces exposed to complex mixtures of PAHs which include BaP. Studies include asphalt works, coke production plants, aluminium smelters and occupations where exposure to coal tar, coal tar pitches and soot occurs. Such studies clearly showed an elevated incidence of lung tumours following inhalation and skin tumours following chronic skin contact. It is difficult to assess with any confidence the contribution of BaP or any other individual PAH to such findings [3, 4, 20].

In the 2012 evaluation, IARC classified BaP as carcinogenic to humans (Group 1). It was concluded that BaP contributes to the genotoxic and carcinogenic effects resulting from occupational exposure to complex PAHs mixtures. The robust animal evidence and consistent and coherent mechanistic evidence from experimental and human studies provide biological plausibility to support the overall classification [19].

Estimated cumulative exposure to BaP of 100 µg/m³ (equivalent to 3.3 µg/m³ for 30 years) in the aluminium smelting industry has been associated with a 2.68 fold increase in the incidence of lung cancer [1].

Reproductive and developmental toxicity

There is evidence to suggest that exposure to PAHs may cause developmental effects in humans. This is supported by evidence in animal developmental studies.

PAH-DNA adducts have been found in fetal cord and maternal blood after maternal exposure to PAHs in ambient air; in light of this the World Health Organisation states that prenatal exposure could increase cancer risk from PAHs [1].

Studies show a dose-response relationship between exposure to PAHs during pregnancy and effects related to intrauterine growth restriction. A study of neonates showed that those with increased levels of PAH-DNA adducts had significantly lower birth weight, length and head circumference [1].

High cord blood levels of BaP-DNA adducts has been associated with decreased birth weight and a reduction in postnatal weight [6].

An association between dietary BaP intake and decreased birth weight, decreased birth length and having a small for gestational age infant was found in women with low vitamin C intake [6].

There is evidence to suggest that BaP may cause developmental neurotoxic effects. Human studies of prenatal environmental PAH exposure (determined by personal air monitoring and measuring BaP-DNA adduct levels in cord blood) have reported neurodevelopmental effects including impaired cognitive ability, impaired neuromuscular function and increased attention problems and anxious/depressed behaviour following prenatal exposure [6]. Infants born close to a coal-fired power station in China had 0.32 ± 0.14 BaP-DNA adducts per 10^8 nucleotides, this level was associated with a decreased motor development at age 2. A 0.1 unit increase in BaP DNA adducts per 10^8 nucleotides, at birth, was associated with a 2 fold greater chance of developmental delay at age 2. After closure of this plant, lower adduct levels were seen in the cord blood of a new cohort and was no longer associated with reduced motor development [1].

Evidence from human studies indicates that PAH or BaP exposure may cause reproductive toxicity in males and females. Studies in adult men exposed to PAH mixtures via occupational exposure or smoking have reported an association between higher levels of BaP-DNA adducts in sperm and male infertility [6, 21]. In a case control study in a Chinese population a strong association was reported between maternal blood BaP-DNA adducts and risk of miscarriage. In a study addressing the probability of conception in women undergoing IVF (in vitro fertilisation), follicular fluid BaP levels were significantly higher in women who did not conceive [6].

Animal and in-vitro data

Inhalation

Rats exposed to BaP dust via inhalation (7.7 mg/m^3 , 2 hours per day, 5 days per week for 4 weeks) showed no treatment related lesions in the lungs or nasal cavities. No dose-response relationship could be demonstrated as only one concentration of BaP was tested [3]. In the same study, kidney sections were also examined and no adverse effects were noted [2, 4]. Similarly, male hamsters did not show any adverse effects following exposure via inhalation to 9.8 mg/m^3 or 44.8 mg/m^3 BaP for 4.5 hours per day, five days per week for 16 weeks [3].

Ingestion

Few data on chronic oral toxicity of BaP in animals are available. Daily oral administration of 120 mg/kg bw BaP to poor affinity AhR mice (DBA/2N) for one to four weeks caused deaths due to myelotoxicity, whereas high affinity mice (C57B1/6N) remained unaffected during the 6 month treatment. Hepatotoxicity, as well as effects on liver and kidney enzymes have also been reported at this concentration [3, 4].

Rats fed $1,100 \text{ mg/kg bw/day}$ BaP in the diet for more than 100 days showed a decreased growth rate [3].

PAHs including BaP have been shown to promote atherosclerotic plaque formation in AhR responsive mice, chickens and pigeons [1].

Dermal Exposure

BaP (16, 32 or 64 g per application) was applied once a week for 29 weeks onto the skin of female mice. Dose-related epidermal thickening and a pronounced inflammatory response of the dermis, amongst other effects were reported in the first weeks of exposure in those administered the high dose, and subsequently in the lower dose groups [2].

Genotoxicity

Several PAHs are mutagenic and genotoxic and induce DNA adducts in vitro and in vivo [1, 11].

BaP has consistently been shown to be positive in in-vitro assays for point mutations in *Salmonella* and for chromosome damage in mammalian cells, in the presence of an exogenous source of metabolic activation. Indeed it is often used as a positive control in such assays [3].

An increase in the same biomarkers of genotoxic effect seen in man on exposure to complex PAH mixtures which may include BaP have also been seen in experimental animals exposed to BaP or anti-BPDE [19]. Such effects include point mutations, sister chromatid exchange, chromosomal aberrations, sperm abnormalities and somatic mutations. BaP induced mutations are notably found in tumour suppressor genes and proto-oncogenes [5].

There is strong evidence that the formation of DNA adducts by BaP is important in mouse lung tumorigenesis and that this mechanism and the formation of radical-cations by BaP is involved in mouse skin carcinogenesis [19].

There is some evidence for the role of BaP in bitumen-fume genotoxicity; in mice exposed to bitumen fume condensates, anti-BPDE–DNA adducts and BaP metabolites have been found in the lungs and urine respectively [7].

Carcinogenicity

IARC concluded that there is sufficient evidence that BaP is carcinogenic to experimental animals [19].

BaP applied directly to the skin of various strains of mice has been reported to induce malignant (and benign) skin tumours, predominantly squamous cell carcinomas. Oral administration of BaP to mice by gavage or diet has yielded an increase in tumour response in lymphoid and hematopoietic tissues and in several organs including the lung, liver oesophagus and tongue. An increase in mammary gland adenocarcinomas has been reported in rats administered BaP by gavage. Lifetime inhalation studies in hamsters gave dose-response related increases in papilloma's and squamous cell-carcinomas in the upper respiratory tract and in the upper digestive tract [19].

Skin tumour development was not observed in AhR deficient mice, however squamous-cell carcinomas of the skin were present in the wild-type mice [19].

Reproductive and developmental toxicity

BaP exposure has been shown to have effects on the development of laboratory animals following prenatal exposure. Developmental effects reported include decreased number of pups, an increase in reabsorptions, reduced pup weight and malformations [1, 2, 6]. Studies in mice exposed to BaP via oral administration during gestation suggest that intrauterine growth restriction, stillbirths and malformations following exposure may be dependent upon the AhR status of the mother and offspring [1-3].

Evidence from animal laboratory studies indicates that gestational exposure to BaP can have an effect on the reproductive function of offspring. Decreases in testis weight, sperm production, testosterone levels and fertility have been reported in male rodents. Effects observed in female mice include decreases in ovary weight, numbers of follicles, corpora lutea and fertility [1, 6].

Persistent neurodevelopmental effects have been observed in rats and mice exposed to BaP via ingestion as neonates; such effects include deficits in learning and memory, anxiety-related behaviours, sensorimotor development and neuromuscular function [6].

Trans-placental exposure to BaP (with dibenzo[a,l]pyrene) has been shown to induce lung and livers tumours in mice [1].

Reproductive effects have been reported in adult laboratory animals exposed to BaP. Reductions in sperm count, motility and production have been reported in various strains of adult male rats and mice and across routes of exposure. Hormonal changes and histological changes in the testis have also been observed in male adult animals. Reduced fertility and reductions in ovary weight have been reported in female adult animals following oral or inhalation exposure to BaP [6].

References

1. World Health Organisation (WHO), WHO guidelines for indoor air quality: selected pollutants. 2010, WHO Regional Office for Europe.
2. Agency for Toxic Substances and Disease Registry (ATSDR), Toxicological Profile for polycyclic aromatic hydrocarbons (PAH). 1995, US department of Health and Human Services: Atlanta, US.
3. International Programme on Chemical Safety (IPCS), Selected non-heterocyclic polycyclic aromatic hydrocarbons. Environmental Health Criteria 202. 1998, WHO: Geneva.
4. Department for Environment Food and Rural Affairs (DEFRA) and Environment Agency (EA), Contaminants in soil: Collation of toxicological data and intake values for humans. Benzo[a]pyrene. , in R&D Publications TOX 2. 2002, Environment Agency: Bristol.
5. International Agency for the Research on Cancer (IARC), Some Non-heterocyclic Polycyclic Aromatic Hydrocarbons and Some Related Exposures, in IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 2010, IARC: Lyon.
6. US EPA, Toxicological Review of Benzo[a]pyrene, I.R.I.S, Editor. 2017: Washington, DC.
7. International Agency for the Research on Cancer (IARC), Bitumens and Bitumen Emissions, and Some N- and S-Heterocyclic Polycyclic Aromatic Hydrocarbons, in IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 2013, IARC: Lyon.
8. European Food Safety Authority (EFSA), Findings of the EFSA Data Collection on Polycyclic Aromatic Hydrocarbons in Food, in A Report from the Unit of Data Collection and Exposure on a Request from the European Commission. 2008.
9. World Health Organisation (WHO), Air quality guidelines for Europe. WHO Regional Publications, European Series, No. 91. 2nd edition. 2000, WHO Regional Office for Europe: Copenhagen.
10. World Health Organisation (WHO), Polynuclear aromatic hydrocarbons in Drinking water. Background document for development of WHO Guidelines for Drinking water quality. 2003, WHO: Geneva.
11. European Food Safety Authority (EFSA), Polycyclic Aromatic Hydrocarbons in Food Scientific Opinion of the Panel on Contaminants in the Food Chain. 2008. The EFSA Journal (2008) 724, 1-114.
12. Food Standards Agency (FSA), Polycyclic Aromatic Hydrocarbons In Cereals, Cereal Products, Vegetables, Vegetable Products And Traditionally Smoked Foods, in Food Survey Information Sheet. 2012.
13. National Physical Laboratory, Annual Report for 2012 on the UK PAH Monitoring and Analysis Network. Report to the Department of Environment, Food and Rural Affairs; the Department of Environment Northern Ireland; the Welsh Government and the Scottish Government. 2014.
14. DEFRA), Air Pollution in the UK 2015. 2016.
15. National Physical Laboratory (NPL), Annual report for 2015 on the UK PAH Monitoring and Analysis Network. 2016.
16. World Health Organisation (WHO), Guidelines for Drinking-Water Quality: Third edition. Vol 1. Recommendations. 1998, WHO: Geneva.
17. Department for Environment Food and Rural Affairs (DEFRA), Technical Guidance Sheet on normal levels of contaminants in English soils: Benzo[a]pyrene. 2012.
18. Department for Environment Food and Rural Affairs (DEFRA), Technical Guidance Sheet on normal levels of contaminants in Welsh soils: Benzo[a]pyrene. 2013.
19. International Agency for the Research on Cancer (IARC), Chemical Agents and Related Occupations, in IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 2012, IARC: Lyon.
20. International Agency for the Research on Cancer (IARC), Polynuclear aromatic compounds, part 1. Chemical, environmental and experimental data. Vol 32, in IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 1983, IARC: Lyon.
21. U.S. Environmental Protection Agency (U.S. EPA), Benzo(a)pyrene (BaP) TEACH Chemical Summary, in U.S. EPA, Toxicity and Exposure Assessment for Children's Health. 2007.

This document from the PHE Centre for Radiation, Chemical and Environmental Hazards reflects understanding and evaluation of the current scientific evidence as presented and referenced here.

First published: August 2018

For queries relating to this document, please contact chemcompendium@phe.gov.uk

For all other enquiries, please contact: phe.enquiries@phe.gov.uk

© Crown copyright 2018, www.gov.uk/phe

Re-use of Crown copyright material (excluding logos) is allowed under the terms of the Open Government Licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ for terms and conditions.


BENZO[a]PYRENE

Benzo[a]pyrene was considered by previous IARC Working Groups in 1972, 1983, and 2005 ([IARC, 1973, 1983, 2010](#)). Since that time new data have become available, which have been incorporated in this *Monograph*, and taken into consideration in the present evaluation.

1. Exposure Data

1.1 Identification of the agent

Chem. Abstr. Services Reg. No.: 50-32-8
Chem. Abstr. Name: Benzo[a]pyrene
IUPAC Systematic Name: Benzo[a]pyrene
Synonyms: BaP; benzo[def]chrysene;
3,4-benzopyrene*; 6,7-benzopyrene*;
benz[a]pyrene; 3,4-benz[a]pyrene*;
3,4-benzpyrene*; 4,5-benzpyrene*
(*alternative numbering conventions)



$C_{20}H_{12}$

Relative molecular mass: 252.31

Description: Yellowish plates, needles from benzene/methanol; crystals may be monoclinic or orthorhombic

Boiling-point: 310–312 °C at 10 mm Hg

Melting-point: 179–179.3 °C; 178.1 °C

Spectroscopy data: Ultraviolet/visual, infrared, fluorescence, mass and nuclear

magnetic-resonance spectral data have been reported

Water solubility: 0.00162 mg/L at 25 °C;
0.0038 mg/L at 25 °C

log K_{ow} (octanol–water): 6.35

Henry's Law Constant: 0.034 Pa m³/mol at 20 °C

From [IARC \(2010\)](#)

1.2 Occurrence and exposure

Benzo[a]pyrene and other polycyclic aromatic hydrocarbons (PAHs) are widespread environmental contaminants formed during incomplete combustion or pyrolysis of organic material. These substances are found in air, water, soils and sediments, generally at trace levels except near their sources. PAHs are present in some foods and in a few pharmaceutical products based on coal tar that are applied to the skin. Tobacco smoke contains high concentrations of PAHs ([IARC, 2010](#)).

1.2.1 Exposure of the general population

The general population can be exposed to benzo[a]pyrene through tobacco smoke, ambient air, water, soils, food and pharmaceutical products. Concentrations of benzo[a]pyrene in

sidestream cigarette smoke have been reported to range from 52 to 95 ng/cigarette — more than three times the concentration in mainstream smoke. Major sources of PAHs in ambient air (both outdoors and indoors) include residential and commercial heating with wood, coal or other biomasses (oil and gas heating produce much lower quantities of PAH), other indoor sources such as cooking and tobacco smoke, and outdoor sources like motor-vehicle exhaust (especially from diesel engines), industrial emissions and forest fires. Average concentrations of individual PAHs in the ambient air in urban areas typically range from 1 to 30 ng/m³; however, concentrations up to several tens of nanograms per cubic metre have been reported in road tunnels, or in large cities that make extensive use of coal or other biomass as residential heating fuel. Estimates of PAH intake from food vary widely, ranging from a few nanograms to a few micrograms per person per day. Sources of PAHs in the diet include barbecued/grilled/broiled and smoke-cured meats; roasted, baked and fried foods (high-temperature processing); bread, cereals and grains (at least in part from gas/flame-drying of grains); and vegetables grown in contaminated soils, or in areas with surface contamination from atmospheric PAH fall-out ([IARC, 2010](#)).

1.2.2 Occupational exposure

Occupational exposure to PAHs occurs primarily through inhalation and via skin contact. Monitoring by means of ambient air-sampling or personal air-sampling at the workplace, to determine individual PAHs, sets of PAHs or surrogates (e.g. coal-tar pitch volatiles) has been used to characterize exposure via inhalation; more recently, biological monitoring methods have been applied to characterize the uptake of certain specific PAHs (e.g. benzo[a]pyrene) to be used as biomarkers of total exposure ([IARC, 2010](#)).

Industries where occupational exposure to benzo[a]pyrene has been measured and reported include: coal liquefaction, coal gasification, coke production and coke ovens, coal-tar distillation, roofing and paving (involving coal-tar pitch), wood impregnation/preservation with creosote, aluminium production (including anode manufacture), carbon-electrode manufacture, chimney sweeping, and power plants. Highest levels of exposure to PAHs are observed in aluminium production (Söderberg process) with values up to 100 µg/m³. Mid-range levels are observed in roofing and paving (e.g. 10–20 µg/m³) and the lowest concentrations (i.e. at or below 1 µg/m³) are observed in coal liquefaction, coal-tar distillation, wood impregnation, chimney sweeping and power plants ([IARC, 2010](#)).

2. Cancer in Humans

No epidemiological data on benzo[a]pyrene alone were available to the Working Group.

3. Cancer in Experimental Animals

Benzo[a]pyrene was considered by three previous Working Groups ([IARC, 1973](#), [1983](#), [2010](#)).

In *IARC Monograph Volume 3* ([IARC, 1973](#)) it was concluded that benzo[a]pyrene produced tumours in all species tested (mouse, rat, hamster, guinea-pig, rabbit, duck, newt, monkey) for which data were reported following exposure by many different routes (oral, dermal, inhalation, intratracheal, intrabronchial, subcutaneous, intraperitoneal, intravenous). Benzo[a]pyrene had both a local and a systemic carcinogenic effect, was an initiator of skin carcinogenesis in mice, and was carcinogenic in single-dose studies and following prenatal and transplacental exposures.

In *IARC Monograph* Volume 32 ([IARC, 1983](#)) no evaluation was made of studies of carcinogenicity in experimental animals published since 1972, but it was concluded that there is *sufficient evidence* for the carcinogenicity of benzo[a]pyrene in experimental animals.

Carcinogenicity studies with administration of benzo[a]pyrene by multiple route of exposure, reported after the initial evaluations, were subsequently reviewed in *IARC Monograph* Volume 92 ([IARC, 2010](#)) and are summarized below ([Table 3.1](#)). See [Table 3.2](#) for an overview of malignant tumours induced in different animal species.

3.1 Skin application

In several studies in which benzo[a]pyrene was applied to the skin of different strains of mice, benign (squamous cell papillomas and keratoacanthomas) and malignant (mainly squamous-cell carcinomas) skin tumours were observed ([Van Duuren et al., 1973](#); [Cavalieri et al., 1977, 1988a](#); [Levin et al., 1977](#); [Habs et al., 1980, 1984](#); [Warshawsky & Barkley, 1987](#); [Albert et al., 1991](#); [Andrews et al., 1991](#); [Warshawsky et al., 1993](#)). No skin-tumour development was seen in *AhR*^{-/-} mice that lacked the aryl hydrocarbon receptor, whereas the heterozygous and wild-type mice developed squamous-cell carcinomas of the skin ([Shimizu et al., 2000](#)).

In a large number of initiation–promotion studies in mice, benzo[a]pyrene was active as an initiator (mainly of squamous-cell papillomas) when applied to the skin ([IARC, 2010](#)).

3.2 Subcutaneous injection

In subcutaneous injection studies of benzo[a]pyrene, malignant tumours (mainly fibrosarcomas) were observed at the injection site in mice ([Kouri et al., 1980](#); [Rippe & Pott, 1989](#)) and rats ([Pott et al., 1973a, b](#); [Rippe & Pott, 1989](#)). No fibrosarcomas were observed in *AhR*^{-/-} mice that

lacked the aryl hydrocarbon receptor, whereas the heterozygous and wild-type mice did develop these tumours ([Shimizu et al., 2000](#)).

In another study, male and female newborn Swiss mice that were given benzo[a]pyrene subcutaneously showed a significant increase in lung-adenoma incidence and multiplicity ([Balansky et al., 2007](#)).

A single study in 12 strains of hamsters resulted in sarcomas at the site of injection in both sexes of all 12 strains ([Homburger et al., 1972](#)).

3.3 Oral administration

After administration of benzo[a]pyrene by gavage or in the diet to mice of different strains ([Spornins et al., 1986](#); [Estensen & Wattenberg, 1993](#); [Weyand et al., 1995](#); [Kroese et al., 1997](#); [Culp et al., 1998](#); [Hakura et al., 1998](#); [Badary et al., 1999](#); [Wijnhoven et al., 2000](#); [Estensen et al., 2004](#)), increased tumour responses were observed in lymphoid and haematopoietic tissues and in several organs, including the lung, forestomach, liver, oesophagus and tongue.

Oral administration of benzo[a]pyrene to *XPA*^{-/-} mice resulted in a significantly higher increase of lymphomas than that observed in similarly treated *XPA*^{+/-} and *XPA*^{+/+} mice ([de Vries et al., 1997](#)). Benzo[a]pyrene given by gavage to *XPA*^{-/-}/*p53*^{+/-} double-transgenic mice induced tumours (mainly splenic lymphomas and forestomach tumours) much earlier and at higher incidences than in similarly treated single transgenic and wild-type counterparts. These cancer-prone *XPA*^{-/-} or *XPA*^{-/-}/*p53*^{+/-} mice also developed a high incidence of tumours (mainly of the forestomach) when fed benzo[a]pyrene in the diet ([van Oostrom et al., 1999](#); [Hoogervorst et al., 2003](#)).

Oral administration of benzo[a]pyrene by gavage to rats resulted in an increased incidence of mammary gland adenocarcinomas ([el-Bayoumy et al., 1995](#)).

Table 3.1 Carcinogenicity studies of benzo[*a*]pyrene in experimental animals

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Skin application				
Mouse, Swiss ICR/ Ha (F) 52 wk Van Duuren <i>et al.</i> (1973)	0 (untreated), 0 (vehicle control), 5 µg/animal, 3 × /wk, 52 wk 50/group	Skin T: 0/50, 0/50, 23/50 (46%; 13 P; 10 C)	+	NR (acetone)
Mouse, Swiss (F) 38–65 wk Cavalieri <i>et al.</i> (1977)	0 and 0.396 µmol [0.1 mg] per animal, twice/wk, 30 wk 40/group	Skin T: 0% [0/29], 78.9% [30/38] (7 P, 7 K, 36 C, 1 malignant Schwannoma)	+	99% (acetone)
Mouse, C57BL/6J (F) 60 wk Levin <i>et al.</i> (1977)	Experiment 1 and 2: 0 (DMSO/ acetone), 0.02 [5.28 µg], 0.1 [26.43 µg], 0.4 [105.75 µg] µmol/ animal, once/2 wk, 60 wk (high dose given in two paintings, 30 min apart) Experiment 3: 0 (acetone/NH ₄ OH), 0.025 [6.6 µg], 0.05 [13.21 µg], 0.1 [26.43 µg] µmol/animal, once/2 wk, 60 wk 30/group	Skin T (mainly SCC): Experiment 1–0%, 0%, 38% (13 T), 100% (44 T) Experiment 2–0%, 4% (1 T), 50% (15 T), 100% (40 T) Experiment 3–0%, 7% (2 T), 59% (20 T), 91% (24 T)	+	NR (DMSO/acetone (1:3) or acetone/NH ₄ OH (1 000:1)) Effective number of animals not clearly specified At most, seven animals/group died prematurely without a skin tumour.
Mouse, NMRI (F) 63–109 wk Habs <i>et al.</i> (1984)	0, 2, 4 µg/animal, twice/wk 20/group	Skin T: 0/20, 9/20 (45%; 2 P, 7 C), 17/20 (85%; 17 C)	+	> 96% (acetone)
Mouse C3H/HeJ (M) 99 wk Warshawsky & Barkley (1987)	0 (untreated), 0 (vehicle control) or 12.5 µg/animal, twice/wk 50/group	Skin T: 0/50, 0/50, 48/50 (96%; 47 C, 1 P)	+	99.5% (acetone)
Mouse, Swiss (F) 42 wk Cavalieri <i>et al.</i> (1988a)	0, 0.1 [26.4 µg], 0.4 [105.7 µg] µmol/ animal, twice/wk, 20 wk 30/group	Skin T incidence: 0/30, 26/29 (90%; SGA, 3 P, 23 SCC), 26/30 (90%; 2 P, 26 SCC)	+	Purified [NR] (acetone)

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Mouse, C3H/fCum (M) Experiment 1: 15 mo Experiment 2: 18 mo Experiment 3: 18 mo Kouri et al. (1980)	Experiment 1: 0 (trioctanoin control), 0 (DMSO control), 0.9 µmol [0.23 mg] in trioctanoin or DMSO, Experiment 2: 0 (trioctanoin control), 0 (DMSO control), 0.9 µmol [0.23 mg] in trioctanoin or in DMSO, Experiment 3: 0 (trioctanoin/ DMSO, 100:1), 0.9 µmol [0.23 mg] in trioctanoin/DMSO (100:1), 1 × 20 or 40/group	FibroS at injection site: <i>Experiment 1</i> – 0/16, 0/20, 15/18 (83%), 12/19 (63%) <i>Experiment 2</i> – 0/20, 0/18, 14/18 (78%), 7/19 (37%) <i>Experiment 3</i> – 0/20, 36/40 (90%)	+	Pure (trioctanoin, DMSO)
Mouse, NR (F) 78 wk Rippe & Pott (1989)	0, 10, 100 µg/ animal, 1 × NR/group	S at injection site: 1/30 (3%), 13/30 (43%), 20/30 (67%)	+	NR (tricaprylin)
Mouse, Swiss (newborn) (M, F) 75–200 d Balansky et al. (2007)	0 and 1.0 mg/animal, 1 × 12–15 M/group, 12–15 F/group	Lung A: M – 0/15, 9/12; F – 0/15, 11/12	$P < 0.001$	Pure (olive oil)
Rat, Wistar (F) ~530 d Pott et al. (1973a)	0, 33, 100, 300, 900, 2 700 µg/ animal, 1 × 50/group	T (mainly fibroS) at injection site [incidence derived from dose–response curves]: 2/50 (~4%), 4/50 (~8%), 7/50 (~14%), 23/50 (~46%), 35/50 (~70%), 38/50 (~76%)	+	NR (tricaprylin)
Rat, NR (F) 132 wk Rippe & Pott (1989)	0 and 1 mg, 1 × NR/group	S at injection site: 0/24 (0%), 20/24 (83%)	+	NR (tricaprylin)
Rat, NR (F) 132 wk Rippe & Pott (1989)	0 and 15 mg, 1 × NR/group	S at injection site: 1/24 (4%), 19/24 (79%)	+	NR (DMSO)

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian, RB (randomly bred), BIO inbred strains designated as: I.5, 4.22, 4.24, 7.88, 12.14, 15.16, 45.5, 54.7, 82.73, 86.93, 87.20 (M, F) 53 wk Homburger et al. (1972)	500 µg/animal, 1 × 25 M/group, 25 F/group	FibroS at injection site: RB–M, 4/25 (16%); F, 6/23 (26%) 1.5–M, 5/25 (20%); F, 4/23 (17%) 4.22–M, 3/25 (12%); F, 8/25 (32%) 4.24–M, not tested; F, 9/25 (36%) 7.88–M, 13/25 (52%); F, 5/23 (23%) 12.14–M, 3/25 (12%); F, 9/22 (41%) 15.16–M, 9/25 (36%); F, 16/25 (64%) 45.5–M, 12/25 (48%); F, 7/15 (47%) 54.7–M, 5/25 (20%); F, 5/25 (20%) 82.73–M, 4/21 (19%); F, 4/24 (17%) 86.93–M, 9/25 (36%); F, 8/25 (32%) 87.20–M, 16/25 (64%); F, 11/25 (42%)	+	NR (tricaprylin) No subcutaneous T observed in historical controls
Oral administration				
Mouse, A/I (F) 260 d Weyand et al. (1995)	0, 16, 98 ppm (total dose; 0, 11, 67 mg) in the diet 30/group	Lung T: 4/21 (19%; 4 A; 0.19 ± 0.09 A/animal), 9/25 (36%*); 7 A, 2 AdC; 0.48 ± 0.14 T/animal), 14/27 (52%*); 14 A; 0.59 ± 0.12 A/animal) Forestomach T: (0%) 0/21, (5/25) (20%; 3 P, 2 C; 0.24 ± 0.11** T/animal), 27/27 (100%*); 13 P, 14 C; 4.22 ± 0.41**)	*P < 0.05 **P < 0.001	NR (gel diet)
Mouse B6C3F1 (F) 2 yr Culp et al. (1998)	0 (acetone control diet), 5 ppm, 25 ppm, 100 ppm in the diet 48/group	Liver (A): 2/48 (4%), 7/48 (15%), 5/47 (11%), 0/45 (0%) Lung (A and/or C): 5/48 (10%), 0/48 (0%), 4/45 (9%), 0/48 (0%) Forestomach (P and/or C): 1/48 (2%), 3/47 (6%), 36/46 (78%****), 46/47 (98%****) Oesophagus (P and/or C): 0/48 (0%), 0/48 (0%), 2/45 (4%), 27/46 (59%**) Tongue (P and/or C): 0/48 (0%), 0/48 (0%), 2/46 (4%), 23/48 (48%****) Larynx (P and/or C): 0/35 (0%), 0/35 (0%), 3/34 (9%), 5/38 (13%*)	*P < 0.014 **P < 0.0014 ***P < 0.0003 ****P < 0.00001	98.5% (acetone)
Mouse, Swiss albino, inbred (F) 27 wk Badary et al. (1999)	0 and 1 mg/animal by gavage, twice/wk, 4 wk 10/group	0, 10/10 (100%) (forestomach P; multiplicity, 7.11 ± 1.05)	+	Highest purity grade (corn oil) Drinking-water contained 0.005% ethanol

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Mouse, CSB ^{-/-} or wild-type (CSP [±] or CSB ^{+/+}) (M, F) 52 wk Wijnhoven et al. (2000)	0 and 13 mg/kg bw by gavage, 3 × /wk, 13 wk 6–18 M/group, 6–13 F/group	Wild-type: 5/27 (14 M, 13 F; 19%; 4 bronchiolo-alveolar A, 2 lymphoma), 17/29* (18 M, 11 F; 59%; 6 bronchiolo-alveolar A, 10 forestomach P, 2 forestomach SCC, 2 histiocytic S, 2 hepatocellular A, 1 intestinal AdC, 1 skin P) CSB ^{-/-} : 0/13 (6 M, 7 F), 7/12** (6 M, 6 F; 58%; 2 bronchiolo-alveolar A, 2 uterine S, 1 forestomach SCC, 1 intestinal AdC, 1 skin histiocytic S)	*P = 0.0023 **P = 0.0017	NR (soya oil)
Rat, Crl:CD(SD)BR (F) 49 wk el-Bayoumy et al. (1995)	0 and 50 µmol/animal, once/wk, 8 wk by gavage 30/group	Mammary T incidence: 11/30 [37%] [incidence not clearly specified] (8 desmoplastic A, 2 A, 1 AdC), 29/30 (96.7%; 8 fibroA**, 17 desmoplastic A*, 7 A, 22 AdC**) Numbers of mammary T: controls, 14 desmoplastic A, 2 A, 1 AdC; treated animals, 14 fibroA*, 35 desmoplastic A, 11 A, 56 AdC**	*P < 0.05 **P < 0.01	99% (trioctanoin)
Intraperitoneal injection				
Mouse, B6C3F ₁ ; C3A/JF ₁ (M, F) 90 wk or lifetime Vesselinovitch et al. (1975a, b)	0, 75, 150 µg in 10 µL/g bw, 1 × at 1, 15, 42 d of age 30–63/group, 96–100 controls/group	B6C3F ₁ mice (all ages combined): Liver T (A and hepatocellular C)– M, 1/98 (1%), 69/162 (43%), 81/165 (49%); F, 0/96 (0%), 7/147 (5%), 10/126 (8%) Lung T (A and AdC)– M, 7/98 (7%), 57/162 (35%), 73/165 (44%); F, 2/90 (2%), 53/147 (36%), 50/126 (40%) Forestomach T (P and SCC)– M, 0/98 (0%), 39/162 (24%), 64/165 (39%); F, 0/96 (0%), 22/147 (15%), 40/126 (32%) Lymphoreticular T (mainly reticulum-cell S)– M, 2/98 (2%), 104/314 (33%) (high- and low-dose groups combined); F, 2/96 (2%), 148/281 (53%) (high- and low-dose groups combined)	+	NR (trioctanoin)

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Vesselinovitch et al. (1975a, b) Contd.			+	
		C3A/JF1 mice (all ages combined): Liver T (A and hepatocellular C)– M, 3/97 (3%), 30/148 (20%), 33/137 (24%); F, 0/100 (0%), 1.3% 2/126 (1.3%), 2/153 1.3%) Lung T (A and AdC)– M, 49/97 (49%), 1 438/148 (93%), 125/137 (91%); F, 26/100 (26%), 115/126 (91%), 141/153 (92%) Forestomach T (P and SCC)– M, 0/97 (0%), 18/148 (12%), 42/137 (31%); F, 0/100 (0%), 18/126 (14%), 31/153 (20%) Lymphoreticular T (mainly reticulum-cell S)– M, 0/97 (0%), 26/285 (9%); F, 2/100 (2%), 50/278 (18%) (high- and low-dose groups combined)		
Mouse, CD-1 (M, F) 1 yr Wislocki et al. (1986)	0 and 560 nmol [148 µg] (total dose; given as 1/7, 2/7, 4/7 on PND 0, 8, 15) 37 M/group, 27 F/group	Liver T: M, 2/28 (7%; 2 A), 18/37* (49%; 11 A, 7 C*); F, no liver T found Lung T: M, 1/28 (4%; 1 A), 13/37** (35%; 13 A); F, 0/31, 13/27** (48%) (13 A) Malignant lymphoma: M, 1/28 (4%), 2/37 (5%); F, 1/31 (3%), 4/27 (15%)	* <i>P</i> < 0.005 ** <i>P</i> < 0.05	> 99% (DMSO)
Mouse, CD-1 (M, F) 52 wk Lavoie et al. (1987)	0 and 1.1 µmol [290 µg] (total dose; given as 1/6, 2/6, 4/6 on PND 1, 8, 15) 17 M/group, 14–18 F/group	Liver T: M, 1/17 (6%; 1 H), 13/17* (76%; 9 hepatic A, 4 H); F, 0/18, 0/14 Lung A: M, 0/17, 14/17* (82%); F, 0/18, 9/14** (64%)	* <i>P</i> < 0.005 **[<i>P</i> < 0.0005]	> 99% (DMSO)
Mouse, Swiss-Webster BLU: Ha(ICR) (M, F) 26 wk Busby et al. (1989)	0 and 59.5 µg (total dose; given as 8.5, 17, 34 µg on PND 1, 8, 15) NR/group	Lung T: M, 12/91 (13%; 12 A, 1 AdC; 0.15 ± 0.04 T/mouse), 13/28 (46%; 13 A; 0.71 ± 0.19 A/mouse); F, 7/101 (7%; 7 A; 0.08 ± 0.03 A/mouse), 18/27 (67%; 18 A, 1 AdC; 1.19 ± 0.21 T/mouse)	+	> 99% (DMSO) statistics NR
Mouse, NR, newborn (M, F) 30 wk Rippe & Pott (1989)	0, 10, 100 µg/animal, 1 × NR/group	Lung T: 13% [5/38] (0.13 T/animal), 16% [5/31] (0.23 T/animal), 64% [21/33] (2.52 T/animal)	+	NR (saline solution + 1% gelatine + 0.4% Tween 20) Type of lung tumour NR; statistics NR
Mouse, A/J (F) 260 d Weyand et al. (1995)	0 (untreated), 0 (vehicle control), 1.79 mg/animal, 1 × 29–30/group	Lung T: 7/30 (23%; 7 A; 0.27 ± 0.12 A/animal), 11/30 (37%; 11 A; 0.43 ± 0.11 A/animal), 29/29* (100%; 27 A, 2 AdC; 15.8 ± 1.28** T/animal); forestomach T: 0/30 (0%), 0/30 (0%), 24/29** (83%; 15 P, 9 C; 1.83 ± 0.25** T/animal)	* <i>P</i> < 0.05 ** <i>P</i> < 0.001	NR (tricaprylin)

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Mouse, B6C3F1 infant (M, F) 26 wk, 39 wk, 52 wk Rodriguez et al. (1997)	0 (untreated), 0 (vehicle controls), 125, 250, 375 µg/7 g bw, 1 × > 30 M/group, > 30 F/group	Liver T (M): At wk 26: 0/41, 0/58, 0/29, 0/25, 3/34 (9%; multiplicity, 1.0); at wk 39: 0/34, 0/59, 6/26 (23%; multiplicity, 1.0), 13/34 (38%); multiplicity, 1.9), 15/23 (65%); multiplicity, 1.9); at wk 52: 4/64 (6%; multiplicity, 1.0), 3/63 (5%); multiplicity, 1.0), 13/29 (45%); multiplicity, 1.8), 14/27 (52%); multiplicity, 2.2), 19/24 (79%); multiplicity, 2.5) No liver T in F	+	NR (corn oil) No forestomach tumours
Mouse, CD-1 (M) 12 mo Von Tungeln et al. (1999)	0, 100, 400 nmol [26, 111 µg]/ animal (total dose; given as 1/7, 2/7, 4/7 on PND 1, 8, 15) 24/group	Liver T: 3/20 (15%); 1 A, 2 C; 1.7 T/liver section), 5/21 (24%); 4 A, 1 C; 1.5 T/liver section), 9/20 (45%; 7 A*, 2 C; > 2.3 T/liver section) Lung T: 4/20 (20%); 4 A; 1.0 T/lung section), 1/21 (5%); 1 A; 1.0 T/lung section), 9/20 (45%); 7 A, 2 C; 1.9 T/lung section)	*P = 0.0234	> 99% (DMSO)
Rat, Wistar (F) ~112 wk Roller et al. (1992)	0 and 5 mg/animal, 1 × NR/group	Abdominal mesothelioma and S: 3/41 (7.3%), 33/37 (89.2%)	+	NR (3:1 mixture of tricaprylin/ beeswax) Limited reporting
Rat, Wistar (F) ~116 wk Roller et al. (1992)	5 mg/animal, 1 × NR/group	Abdominal mesothelioma and S: 19/38 (50%); historical controls, 11/369 (3%)	+	NR (saline solution) No control; limited reporting of tumour data
Inhalation				
Hamster, Syrian golden (M) Lifetime Thyssen et al. (1981)	0, 2.2, 9.5, 46.5 mg/m ³ , 4.5 h/d, 7 d/ wk, 10 wk; thereafter 3 h/d, 7 d/wk (total average doses: 0, 29, 127, 383 mg/animal) 24/group (+ animals added during the study)	Respiratory tract T: (polyps, P, SCC) – 0/27, 0/27, 34.6% [9/26; 3 nasal, 8 laryngeal, 1 tracheal], 52% [13/25; 1 nasal, 13 laryngeal, 3 tracheal; no bronchogenic T] Upper digestive tract T: (polyps, P, SCC) – 0/27, 0/27, 26.9% [6/26; 6 pharyngeal, 1 forestomach], 56% [14/25; 14 pharyngeal, 2 oesophageal, 1 forestomach]	+	NR (0.1% saline solution); particle size, > 99% diameter 0.2–0.5 µm, > 80% diameter 0.2–0.3 µm Survival decreased for high dose-exposed animals (59 wk) vs other groups (96 wk).

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Intrapulmonary injection				
Rat, OM (F) 64 (high-dose group)–133 wk (untreated controls) Deutsch-Wenzel et al. (1983)	0 (untreated), 0 (vehicle control), 0.1, 0.3, 1.0 mg/animal, 1 × 35/group	Lung T: [0/35] (0%), [0/35] (0%), [10/35] (28.6%) (4 epidermoid C; 6 pleomorphic S), [23/35] (65.7%) (21 epidermoid C; 2 pleomorphic S), [33/35] (94.3%) (33 epidermoid C)	+	99.1% (1:1 mixture of beeswax and tricaprylin)
Rats, F344/NSlc (M) 104 wk Iwagawa et al. (1989)	0, 0.03, 0.1, 0.3, 1.0 mg/ animal, 1 × NR/group	Lung T: 0/40, 1/29 (3%; 1 undifferentiated T), 7/30 (23%; 6 SCC, 1 undifferentiated T), 22/29 (76%; 20 SCC, 2 undifferentiated T), 9/13 (69%; 9 SCC)	+	NR (1:1 mixture of beeswax/ tricaprylin)
Rat, Osborne-Mendel (F) 134 wk (low-dose group)–140 wk (vehicle controls) Wenzel-Hartung et al. (1990)	0 (untreated), 0 (vehicle control), 30, 100, 300 µg/ animal, 1 × 35/group	Lung T: [0/35] (0%), [0/35] (0%), [3/35] (8.6%; 3 SCC), [11/35] (31.4%; 11 SCC), [27/35] (77.1%; 27 SCC).	+	99.6% (beeswax/trioctanoin mixture of varying composition) SCC predominantly keratinized
Rat, F344/DuCrj (M) 100 wk Horikawa et al. (1991)	0, 50, 100, 200 µg/animal, 1 × 9–10/group	Lung T: 0/19, 0/10, 3/10 (30%; 2 SCC, 1 AdSC), 4/9 (44.4%; 3 SCC, 1 undifferentiated T)	+	NR (1:1 mixture of beeswax/ tricaprylin)
Intratracheal administration				
Rat, Wistar-WU/ Kisslegg (F) 124–126 wk Pott et al. (1987)	0 and 1 mg/animal, once/wk, 20 wk NR/group	Lung T: 0/40, 7/36 (19%; 1 A, 5 SCC, 1 mixed AdC/SCC)	+	NR (0.9% saline solution)
Rat, Sprague-Dawley (M, F) Controls, 131 wk; treated animals, 112 wk Steinhoff et al. (1991)	0 and 0 (physiological saline), 7 mg/kg bw/instillation (physiological saline with Tween 60), once/2 wk, 44 wk 20 or 50/group	M: 0/50, 0/50, 19/20 (95%; 19 malignant lung T) F: 0/50, 0/50, 19/20 (95%; 18 malignant, 1 benign lung T)	+	NR (physiological saline solution with or without Tween 60) Limited histology

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (M, F) 78 wk Feron (1972)	0 (only for M), 1 mg/animal, once/ wk, 36 wk 35/group	Respiratory tract T/adenomatoid lesions: M–6/27 (22%); 1 tracheal P, 5 pulmonary adenomatoid lesion, 19/29 (66%); 1 tracheal P, 17 SCC, 26 pulmonary adenomatoid lesion, 5 A, 1 AdC, 1 SCC F–22/27 (81%); 1 laryngeal SCC, 16 tracheal SCC, 2 bronchial A, 1 AdC, 21 pulmonary adenomatoid lesion, 8 A, 1 AdC)	+	> 99% (0.9% saline solution) No female controls; Statistics NR
Hamster, Syrian golden (M) 78 wk Feron et al. (1973)	0, 0.0625, 0.125, 0.25, 0.5, 1.0 mg/ animal, once/wk, 52 wk 30/group	Respiratory tract T: 0/29, 3/30 (10%); 3 tracheal P, 1 pulmonary A), 4/30 (13%); 1 tracheal P, 4 pulmonary A), 9/30 (30%); 5 tracheal P, 7 pulmonary A), 25/29 (86%); 2 tracheal polyp, 9 P, 5 SCC, 1 AdSC, 1 fibroS, 2 bronchial polyp, 1 P, 2 SCC, 1 AdSC), 26/28 (93%); 6 tracheal P, 11 SCC, 1 AdSC, 1 bronchial polyp, 2 P, 4 SCC, 2 AdSC, 4 AdC, 1 anaplastic C, 16 pulmonary A, 4 SCC, 3 AdSC, 1 AdC, 2 anaplastic C)	+	NR (0.9% saline solution)
Hamster, Syrian golden (M, F) M, 67–88 wk; F, 60–88 wk Henry et al. (1973)	0, 13.3–15.5 mg/animal, once/wk, 8 wk 50/group, 25 controls/group	Respiratory tract T: Controls– 1 tracheal polyp, 6 pulmonary bronchiolar adenomatoid lesions/47 animals Treated animals– 26/65 (40%); 1 nasal polyp; 6 laryngeal polyps, 1 P, 1 A, 1 AdC, 7 tracheal polyps, 1 AdC, 1 SCC, 1 fibroS, 2 bronchial AdC, 13 pulmonary bronchiolar adenomatoid lesion, 3 A, 5 AdC, 1 SCC, 2 anaplastic C, 1 mixed C, 1 myelogenous leukaemia, 1 neurofibroS) T at other sites: Controls– 1 renal A Treated animals– 3 blast-cell leukaemia, 2 adrenocortical A, 1 renal AdC, 1 oesophageal fibroS, 1 haemangioma	+	NR (0.5% gelatine in 0.9% saline solution) Tumour data for M and F combined; statistics NR

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (M, F) 60 wk Kobayashi (1975)	0 and 1 mg/animal, once/wk, 30 wk 20–32 M/group, 20–28 F/group	Respiratory tract T: M–0/20, 11/26 (42.3%); 1 laryngeal polyp, 1 tracheal polyp, 1 P, 1 bronchial SCC, 9 lung A, 7 AdC, 3 SCC, 1 anaplastic C, 2 AdSC) F–0/20, 14/26 (53.8%); 1 laryngeal P, 2 tracheal polyps, 1 bronchial SCC, 10 lung A, 3 AdC, 1 SCC)	+	NR (0.9% saline)
Hamster, Syrian golden (M, F) 78 wk Kruysse & Feron (1976)	0 (untreated), 0 (vehicle controls), 1 mg/animal, once/2 wk, 52 wk 17 or 40/group	Respiratory tract T: M–0/40, 0/40, 13/14 (93%); 2 laryngeal P, 1 SCC, 4 tracheal P, 3 SCC, 1 anaplastic C, 1 S, 1 bronchial SCC, 1 AdC, 5 pulmonary A, 1 AdC) F–0/40, 0/40, 7/12 (58%); 2 tracheal P, 3 SCC, 1 bronchial P, 5 pulmonary A)	+	> 99% (saline solution)
Hamster, Syrian golden (M) 100 wk Sellakumar <i>et al.</i> (1976)	0 (untreated), 3 mg/animal, once/ wk, 10 wk 48/group	Respiratory tract T: 0/48, 7/48 (15%); 2 laryngeal P, 4 tracheal P, 1 lung A) T at other sites: 6/48 (13%); 3 forestomach P, 2 lymphoma, 1 anaplastic C), 26/48 (54%); 21 forestomach P, 1 skin melanoma, 1 liver haemangioma, 1 adrenocorticoA, 3 adrenocorticoC)	+	> 99% (0.9% saline solution)
Hamster, Syrian golden (M, F) Experiment 1: up to 89 wk for M and 70 wk for F Experiment 2: up to 83 wk for M and 68 wk for F Ketkar <i>et al.</i> (1977)	Experiment 1: 0, 4, 8, 16 mg in 0.9% saline solution/animal, 1 × 30/group Experiment 2: 0, 4, 8, 16 mg in Tris buffer/ animal, 1 × 30/group	Respiratory tract T: Experiment 1– M 0/24, 3/30 (10%); 1 laryngeal P, 1 tracheal P, 1 lung S), 5/28 (18%); 1 laryngeal SCC, 1 tracheal P, 4 lung S), 4/27 (15%); 3 tracheal P, 1 lung A, 1 S) F 0/28, 3/29 (10%); 1 tracheal P, 2 lung A), 1/30 (3%); 1 lung A), 3/28 (13%); 1 laryngeal P, 2 lung A) Experiment 2– M 0/27, 5/24 (21%); 1 tracheal P, 5 lung A), 13/25 (52%); 1 laryngeal P, 7 tracheal P, 4 lung A, 3 AdC), 8/27 (30%); 2 laryngeal P, 1 SCC, 3 tracheal P, 3 lung A) F 0/27, 3/27 (11%); 2 tracheal P, 1 lung AdC), 2/29 (7%); 2 tracheal P), 8/29 (28%); 1 laryngeal P, 4 tracheal P, 5 lung A)	+	97% (0.9% saline solution or Tris buffer)

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (M, F) 81 wk Feron & Kruysse (1978)	0 (untreated), 0 (vehicle controls), 0.35, 0.7 mg/animal, once/wk, 52 wk 15 or 30/group	Respiratory tract T: M-0/30 (untreated and vehicle controls combined), 4/29 (14%; 2 tracheal P, 1 bronchial P, 2 pulmonary A), 19/30 (63%; 1 laryngeal P, 5 tracheal P, 1 SCC, 1 anaplastic C, 1 S, 2 bronchial P, 1 AdC, 11 pulmonary A, 2 AdC, 1 SCC, 1 anaplastic C) F-0/28 (untreated and vehicle controls combined), 3/27 (11%; 1 laryngeal P, 1 bronchial P, 1 pulmonary A), 7/24 (29%; 1 tracheal P, 2 SCC, 1 bronchial AdC, 5 pulmonary A)	+	> 99% (0.9% saline solution) Statistics NR
Hamster, Syrian golden (M, F) Average survival up to 41 wk for M and 35 wk for F Ketkar et al. (1978)	0, 0.1, 0.33, 1.0 mg/animal, once/wk 30/group	Respiratory tract T: M-0/29, 5/26 (19%; 5 bronchiogenic A), 7/29 (24%; 5 tracheal P, 2 bronchiogenic A), 6/27 (22%; 5 tracheal P, 2 bronchiogenic A) F-0/30, 12/30 (40%; 1 tracheal P, 1 SCC, 10 bronchiogenic A), 10/28 (36%; 7 tracheal P, 5 bronchiogenic A, 1 SCC), 6/30 (20%; 3 tracheal P, 3 bronchiogenic A, 3 SCC)	+	97% (10% bovine serum albumin) Average survival time much lower in the high-dose group than in the other groups
Hamster, Syrian golden (M, F) Lifetime, up to 90 wk Stenbäck & Rowland (1978)	0, 3 mg large particles, 3 mg small particles/animal, once/wk, 18 wk 48 (M + F)/group	Respiratory tract T (M + F combined): 0/46, 31/47 (66%; 5 laryngeal P, 12 tracheal P, 20 SCC, 2 unspecified T, 2 bronchial P, 9 SCC, 3 A, 2 anaplastic C), 5/46 (11%; 1 laryngeal P, 1 SCC, 4 tracheal P)	+	99.4% (0.9% saline solution); particle size by weight: large-98% < 30 µm, 90% < 20 µm, 36% < 10 µm, 10% < 5 µm; small-98% < 10 µm, 79% < 5 µm, 5% < 1 µm
Hamster, Syrian golden (M) Average survival up to 88 wk Ketkar et al. (1979)	0 (untreated), 0 (vehicle controls), 0.125, 0.25, 0.5, 1.0 mg/animal, once/wk 30/group	Respiratory tract T: 0/29, 0/28, 9/29 (31%; 2 laryngeal polyps/P, 1 tracheal P, 1 SCC, 2 lung A, 2 SCC, 5 AdC), 24/29 (83%; 1 nasal SCC, 2 laryngeal polyps/P, 4 tracheal P, 9 SCC, 5 lung A, 5 SCC, 11 AdC), 19/29 (66%; 1 laryngeal P, 2 SCC, 5 tracheal P, 11 SCC, 7 lung SCC, 2 AdC), 9/29 (31%; 1 laryngeal P, 1 SCC, 1 tracheal P, 5 SCC, 1 lung A, 4 SCC)	P < 0.001, all treated groups	97% (Tris buffer + 0.9% saline solution); particle size: majority < 10 µm but particles up to 80 µm also present Average survival in two highest-dose groups much lower than that in the other groups due to many early deaths from pulmonary lesions other than tumours

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (M, F) 105 wk Feron et al. (1980)	0 (untreated), 0 (gelatine in 0.9% saline), 0.5, 1.0 mg fine particles, 0.5, 1.0 mg coarse particles, 1.0 mg wide-range particles/animal, once/wk, 52 wk 30–35/group	Respiratory tract T: M–0/29, 2/34 (6%); 2 laryngeal P, 7/34 (21%); 1 laryngeal P, 6 tracheal P, 1 lung A, 6/31 (19%); 2 laryngeal P, 1 tracheal P, 1 S, 1 pulmonary A, 13/31 (42%); 2 laryngeal P, 3 tracheal P, 9 pulmonary A, 25/34 (74%); 2 laryngeal P, 9 tracheal P, 4 SCC, 2 S, 1 pulmonary A, 1 AdC), 23/34 (68%); 2 laryngeal P, 1 SCC, 6 tracheal P, 2 SCC, 1 bronchial P, 1 SCC, 13 pulmonary A, 2 AdC, 2 anaplastic C) F–0/28, 2/33 (6%); 1 tracheal P, 1 pulmonary A, 2/34 (6%); 1 bronchial P, 1 A, 5/32 (16%); 1 laryngeal P, 2 tracheal P, 3 pulmonary A, 9/32 (28%); 2 laryngeal P, 5 tracheal P, 6 pulmonary A, 19/32 (31%); 4 tracheal P, 1 SCC, 1 S, 1 bronchial P, 7 pulmonary A, 1 AdC), 11/34 (34%); 1 laryngeal P, 3 tracheal P, 2 bronchial P, 7 pulmonary A, 1 AdC)	+	NR; particles size by weight: fine, 77% < 5.2 µm, 60% < 3.9 µm; coarse, 77% < 42 µm, 3% < 16 µm; wide-range, 72% < 30 µm, 19% < 10 µm (gelatine in 0.9% saline solution) Statistics NR
Hamster, Syrian golden (M) 129 wk Godleski et al. (1984)	0 and 5 mg/animal, once/wk, 15 wk 80/group	Malignant T: 4/80 (5%); 1 multicentric undifferentiated lung C, 3 lymphoma), 25/80* (31%); 9 SCC, 2 undifferentiated C of the respiratory tract, 5 lymphoma, 1 SCC, 2 AdC of the gastrointestinal tract, 2 soft-tissue T, 1 hepatoma, 2 mouth SCC, 1 skin C)	*P < 0.001	> 99% (0.5% gelatine in 0.9% saline solution)
Intratracheal administration of combinations of benzo[a]pyrene and 'particles/fibres'				
Rat, Sprague-Dawley (M, F) Up to 130 wk Steinhoff et al. (1991)	0, (untreated), 0 (physiological saline), 10–40 mg/kg bw Bayferrox 130 (96.2% cubic α -Fe ₂ O ₃), 10–40 mg/kg bw Bayferrox 920 (86.1% fibrous α -FeOOH), 7 mg/kg bw, 7 mg/kg bw + 10–40 mg/kg bw Bayferrox 130, 7 mg/kg bw + 10–40 mg/kg bw Bayferrox 920, ~once/2 wk, 44–~130 wk 20 or 50/group	Lung T: M–0/50, 0/50, 0/50, 0/50, 19 malignant T in 20 animals, 21 malignant and 1 benign T in 20 animals, 17 malignant and 1 benign T in 20 animals F–0/50, 0/50, 0/50, 1 malignant and 1 benign T in 50 animals, 18 malignant and 1 benign T in 20 animals, 16 malignant T in 20 animals, 17 malignant and 2 benign T in 20 animals	+	NR (physiological saline solution with or without Tween 60); Bayferrox 130, Bayferrox 920 Limited histology

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (M, F) Lifetime (up to 140 wk) Saffiotti et al. (1972)	Experiment 1 0, 50 mg ferric oxide, 5 mg + 45 mg ferric oxide, 12.5 mg + 37.5 mg ferric oxide/ animal, 1 × Experiment 2 (2 groups/dose level) 5 mg + 5 mg ferric oxide, 10 mg + 10 mg ferric oxide, 15 mg + 15 mg ferric oxide, once/wk, 15 wk 23–110 M/group, 18–107 F/group	Experiment 1: Respiratory tract T– M 0/45, 0/101, 3/92 (3%; 1 tracheal polyp, 1 P, 1 bronchial A), 3/27 (11%; 1 bronchial A, 1 bronchogenic SCC, 1 anaplastic C) F 0/44, 0/89, 4/97 (4%; 1 tracheal polyp, 1 P, 1 bronchiolar A, 1 AdC), 6/33 (18%; 1 bronchial P, 1 A, 2 bronchogenic SCC, 1 anaplastic C, 2 bronchiolar A) Forestomach P– M 5/45 (11%; 6 T), 5/101 (5%; 5 T), 15/92 (16%; 35 T), 8/27 (30%; 16 T) F 2/44 (5%; 2 T), 2/89 (2%; 3 T), 5/97 (5%; 5 T), 4/33 (12%; 6 T) Experiment 2: Respiratory tract T (M + F combined)– 7/50 (14%; 2 tracheal P, 1 SCC, 1 bronchial P, 1 A, 2 SCC, 1 anaplastic C, 1 pulmonary SCC), 8/58 (14%; 2 tracheal polyps, 1 bronchial polyp, 2 SCC, 2 AdC, 2 pulmonary A, 2 AdC), 17/61 (28%; 2 tracheal polyps, 2 P, 5 SCC, 5 bronchial SCC, 1 pulmonary A, 1 SCC, 1 AdC, 1 anaplastic C), 25/60 (42%; 4 tracheal polyps, 3 P, 3 SCC, 4 anaplastic C, 1 bronchial P, 1 SCC, 2 anaplastic C, 4 AdC, 1 A, 2 pulmonary SCC, 2 anaplastic C, 6 A), 25/39 (64%; 1 tracheal P, 10 SCC, 1 anaplastic C, 3 bronchial P, 7 SCC, 11 anaplastic C, 2 AdC, 2 pulmonary SCC, 2 A), 35/55 (64%; 2 laryngeal SCC, 11 tracheal P, 1 polyp, 12 SCC, 1 carcinoS, 2 fibroS, 16 bronchial SCC, 10 anaplastic C, 6 AdC, 3 A, 2 pulmonary A) Forestomach T– M 8/22 (36%; 13 P, 1 SCC), 6/28 (21%; 9 P), 11/34 (32%; 28 P, 1 SCC), 11/30 (37%; 18 P), 5/22 (23%; 10 P), 1/28 (4%; 1 P) F 9/28 (32%; 14 P), 6/30 (20%; 9 P), 5/27 (19%; 20 P), 8/30 (27%; 10 P, 1 SCC), 5/17 (29%; 11 P), 3/27 (11%; 3 P)	+	NR (0.9% saline solution); ferric oxide

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (F) Presumably lifetime Pott et al. (1973b)	340 µg in tricaprylin, the larynx, trachea or bronchi): 340 µg in Tween 60/saline solution, 340 µg in Tween 60/saline solution + 850 µg atmospheric dust/animal, 45 × within a period of 6.5 mo (total dose, ~15 mg; dust, 38 mg) 48/group	Respiratory tract T (benign and malignant T of the larynx, trachea or bronchi): 2/48 (4%), 14/48 (29%), 16/48 (33%)	+	NR (tricaprylin, Tween 60/ saline solution); atmospheric dust from Bochum, Germany (particle size < 5 µm)
Hamster, Syrian (M, F) 100 wk Sellakumar et al. (1973)	0 (untreated), 3 mg + 3 mg ferric oxide, 3 mg + 6 mg ferric oxide, 3 mg + 9 mg ferric oxide, once/2 wk, 20 wk 36/group, 193 controls/group	Respiratory tract T (M + F combined): 0/193, 26/67 (39%); 3 laryngeal polyp, 3 P, 3 SCC, 7 tracheal polyp, 6 P, 2 SCC, 2 bronchial polyp, 5 SCC, 9 AdC, 1 anaplastic C, 7 lung A, 1 AdC), 28/64 (44%); 1 laryngeal polyp, 3 P, 6 SCC, 3 tracheal polyp, 9 P, 3 SCC, 3 bronchial polyp, 1 P, 4 SCC, 3 AdC, 1 anaplastic C, 7 lung A, 4 AdC), 26/66 (39%); 3 laryngeal polyp, 6 SCC, 6 tracheal polyp, 11 P, 1 SCC, 1 bronchial polyp, 1 P, 4 SCC, 4 AdC, 2 anaplastic C, 6 lung A, 6 AdC) Forestomach T: M–0/193 (M + F), 17/32 (53%; 37 P), 10/31 (32%; 16 P, 1 SCC), 6/35 (17%; 15 P) F–0/193 (M + F), 10/35 (29%; 30 P), 12/33 (36%; 25 P), 15/31 (48%; 33 P)	+	NR (0.9% saline solution); ferric oxide
Hamster, Syrian golden (M, F) Lifetime (up to 120 wk) Stenbäck et al. (1975)	0 (untreated), 2 mg + 1 mg magnesium oxide/ animal, once/wk, 20 wk, 3 mg + 3 mg ferric oxide/animal, once/wk, 15 wk 48 or 90/group	Respiratory tract tumours (M + F combined): 0/89, 32/45 [71%] (11 laryngeal P, 3 SCC, 1 tracheal polyp, 20 P, 5 SCC, 1 AdC, 1 bronchial P, 3 A, 8 AdC, 9 SCC, 1 AdSC), 31/44 (70%; 10 laryngeal P, 4 SCC, 8 tracheal P, 12 SCC, 2 anaplastic C, 2 bronchial P, 4 A, 2 AdC, 17 SCC, 3 anaplastic C)	+	NR (0.2% saline solution); ferric oxide, magnesium oxide

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Hamster, Syrian golden (M, F) Lifetime (up to 100 wk) Stenbäck & Rowland (1979)	0 (untreated), 0 (saline), 0 (gelatine in saline), 3 mg silicon dioxide in saline, 1.5 mg manganese dioxide in saline, 3 mg in saline, 3 mg in gelatine/saline, 3 mg + 3 mg silicon dioxide in saline, 1.5 mg + 1.5 mg manganese dioxide in saline/ animal, once/wk, 20 wk 50/group	All T (M + F combined): 2/100 (2%); 2 lymphoma), 1/48 (2%; 2 forestomach P), 2/45 (4%; 2 lymphoma), 0/48 (0%); 2/48 (4%; 1 forestomach P, 1 lymphoma), 18/46 (39%; 1 laryngeal P, 1 SCC, 4 tracheal P, 15 forestomach P), 11/47 (23%; 2 tracheal P, 1 SCC, 3 bronchial SCC, 1 splenic haemangioma, 1 adrenal cortical A, 1 lymphoma, 2 forestomach SCC), 25/48 (52%; 1 laryngeal SCC, 8 tracheal P, 2 SCC, 3 bronchial SCC, 6 lung A, 3 AdC, 10 forestomach P, 1 thyroid A, 1 uterine fibroma, 1 A, 1 lymphoma), 20/48 (42%; 1 laryngeal P, 3 tracheal P, 1 SCC, 1 bronchial SCC, 24 forestomach P, 1 ovarian fibroma, 1 thyroid A, 2 forestomach SCC, 1 squamous-cell fibroma)	+	> 99% (saline, 0.5% gelatine in saline); manganese dioxide, silicon dioxide
Hamster, Syrian golden (M, F) 82 wk Reynders et al. (1985)	0 and 8 mg + 6 mg ferric oxide/ animal, once/wk, 6 wk 35/group	Respiratory tract T: M-0/32, 12/24 (50%); 15 T: 3 laryngeal P, 1 tracheal P, 1 SCC, 2 bronchial polyp, 2 SCC, 1 AdC, 3 pulmonary SCC, 1 AdSC, 1 AdC) F-0/35, 9/26 (35%); 12 T: 1 laryngeal P, 5 tracheal P, 2 bronchial polyp, 2 pulmonary SCC, 1 AdSC, 1 AdC)	+	NR (0.9% saline solution); ferric oxide
Buccal pouch				
Hamster, Syrian golden (M) Up to 40–44 wk with interim kills after 5, 20, and 24–32 wk Solt et al. (1987)	Painting of both buccal pouch with 0, 20 mM solution/animal, twice/ wk, 20 wk 28/group, 20 controls/group	Forestomach P: 0/6, 8/10* (after 40–44 wk) Buccal pouch SCC: 0/6, 1/10 (after 40–44 wk)	*[P < 0.01]	NR (paraffin oil)
Intramammary or intramamillary administration				
Rat, Sprague-Dawley (F) 20 wk Cavalieri et al. (1988a)	0 and 0 (untreated contralateral mammary gland), 4 [1 mg], 16 µmol [4.2 mg] (5 th right mammary gland), 1 × 20/group	Mammary gland T: [0/20] (0%), [0/20] (0%), [10/20] (50%; 6 AdC, 4 fibroS), [16/20] (80%; 8 AdC, 2 fibroA, 10 fibroS)	+	> 99% (no vehicle)

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Rat, Sprague-Dawley (F) 45 wk Cavalieri et al. (1988a, b)	0 and 4 µmol [1 mg]/mammary gland (2nd, 3rd, 4th and 5th mammary gland on both sides injected), 1 × 20/group	Epithelial mammary T: [3/20] (15%; 3 fibroA), [14/20] (70%; 13 AdC, 3 fibroA); multiplicity: controls, 3/3 [1]; treated rats: AdC, 18/13 [1.4]; fibroA, 4/3 [1.3] Mesenchymal (mammary) T: [0/20] (0%), [11/20] (55%; 11 fibroS; multiplicity, 20/11 [1.8]) Skin T: [0/20] (0%), [9/20] (45%; 9 SCC; multiplicity, 11/9 [1.2])	+	> 99% (trioctanoin)
Rat, Sprague-Dawley (F) 24 wk Cavalieri et al. (1991)	0, 0.25 [66 µg], 1 µmol [264 µg]/mammary gland (the 2nd, 3rd, 4th and 5th on both sides), 1 × 20/group	Epithelial mammary gland T: 1/18 (6%; 1 fibroA), 1/20 (5%; 1 AdC), 0/20 (0%) Mesenchymal (mammary) T: 0/18 (0%), 6/20 (30%; 6 fibroS; multiplicity, 7/6), 8/20 (40%; 8 fibroS; multiplicity, 10/8) Skin T: 0/18 (0%), 0/20 (0%), 1/20 (5%; 1 SCC)	+	> 99% (trioctanoin) Statistics NR
Intracolonic instillation				
Mouse, Swiss albino (M, F) 120 wk Toth (1980)	0, 200, 2000 µg/g bw (total doses); control and high-dose group, 10 × /wk instillations of 0 and 200 µg, respectively; low-dose group, 1 instillation 50/group/sex	Malignant lymphoma: M–0/50, 6/50* (12%; 1 histiocytic, 4 lymphocytic, 1 mixed), 7/50** (14%; 2 histiocytic, 3 lymphocytic, 2 mixed) F–11/49 (22%; 5 histiocytic, 6 lymphocytic), 21/50*** (42%; 5 histiocytic, 16 lymphocytic), 18/49 (36%; 6 histiocytic, 8 lymphocytic, 4 mixed) Oesophagus T: M–no tumour F–0/49, 0/50, 5/49 (10%) Forestomach T: M–0/50, 2/50 (4%; 2 P), 10/50**** (20%; 9 P, 1 SCC) F–1/49 (2%; 1 SCC), 5/10 (20%; 3 P, 2 SCC), 11/49**** (22%; 9 P, 2 SCC)	*P < 0.04 **P < 0.02 ***P < 0.053 ****P < 0.006 *****P < 0.0001	98% (olive oil) Anal and skin tumours probably due to release of benzo[a]pyrene through the anal orifice

Table 3.1 (continued)

Species, strain (sex) Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Result or significance	Purity (vehicle) Comments
Toth (1980) Contd.		Anal T : M-0/50, 0/50, 7/50** (14%; 4 P, 3 SCC) F-0/49, 1/50 (2%; 1 P), 6/49* (12%; 1 P, 4 SCC, 1 K) Skin T : M-1/50 (2%; 1 K), 0/50, 13/50***** (26%; 5 P, 7 SCC, 1 K) F-0/49, 2/50 (4%; 2 SCC), 11/49***** (22%; 4 P, 5 SCC, 2 K)		
Mouse, C57Bl/6 (F) 18 mo Anderson et al. (1983)	0 (untreated, olive oil or β -naphthoflavone in olive oil), 1 mg/animal (in olive oil), once/ wk, 14 wk 45–60/group	Forestomach P: 7/34 (21%; multiplicity, 1.1 ± 0.4), 17/18* (94%; multiplicity, $3.2 \pm 2.3^*$) Peritoneal S: 0/40, 5/32* Lymphoma: 1/40 (2.5%), 9/32* (28%)	* $P < 0.05$	99% (olive oil, enzyme inducer β -naphthoflavone) No colon tumours found
Intravaginal application				
Mouse, C57Bl (F) 5 mo Näslund et al. (1987)	Cotton swab soaked in acetone (controls) or 1% solution of benzo[a]pyrene in acetone, twice/ wk 10 or 76/group	0/10, 17/76 (22%; invasive cervical C)	+	NR (acetone)
Intrafetal injection				
Mouse, Swiss (M, F) 12 wk Rossi et al. (1983)	0, 0.4, 4.0, 9.9, 19.8 nmol [0, 0.1, 1, 2.6, 5.2 μ g]/animal, 1 $\times$ 43–56/group	Lung A (M + F combined): 0/37, 1/39 (3%), 10/42 (25%), 10/38 (26%), 12/31 (39%)	+	> 99% (trioctanoin-acetone mixture (1:1))

A, adenoma; AdC, adenocarcinoma; AdSC, adenosquamous carcinoma; bw, body weight; C, carcinoma; d, day or days; DMSO, dimethyl sulfoxide; F, female; H, hepatoma; K, keratocanthoma; M, male; min, minute or minutes; mo, month or months; NH_4OH , ammonium hydroxide; NR, not reported; P, papilloma; PND, postnatal day; S, sarcoma; SCC, squamous-cell carcinoma; SGA, sebaceous gland adenoma; T, tumour; vs, versus; wk, week or weeks; yr, year or years

Table 3.2 Summary of reports of malignant tumours clearly induced in experimental animals by benzo[a]pyrene

Organ site/ species	Lung	Trachea	Larynx	Forestomach	Liver	Lymphoid tissue (lymphoma)	Sarcoma (injection site)	Skin	Mammary gland
Mouse	x			x	x	x	x	x	
Rat	x						x		x
Hamster	x	x	x	x			x		

3.4 Intraperitoneal injection

In a series of studies in newborn and adult mice, intraperitoneal injection of benzo[a]pyrene increased the incidence of liver (adenomas and carcinomas) and lung (adenomas and adenocarcinomas) tumours and, occasionally, forestomach (squamous cell papillomas and carcinomas) and lymphoreticular tumours ([Vesselinovitch et al., 1975a, b](#); [Wislocki et al., 1986](#); [Lavoie et al., 1987](#); [Busby et al., 1989](#); [Rippe & Pott, 1989](#); [Mass et al., 1993](#); [Nesnow et al., 1995](#); [Ross et al., 1995](#); [Weyand et al., 1995](#); [Rodriguez et al., 1997](#); [Von Tungeln et al., 1999](#)).

In one study in rats with a single intraperitoneal injection of benzo[a]pyrene, a high incidence of abdominal mesotheliomas and sarcomas was observed ([Roller et al., 1992](#)).

3.5 Inhalation

In a lifetime inhalation study ([Thyssen et al., 1981](#)) in male hamsters, benzo[a]pyrene induced dose-related increases in the incidence of papillomas and squamous-cell carcinomas in both the upper respiratory tract (nose, larynx and trachea) and the upper digestive tract (pharynx, oesophagus and forestomach).

3.6 Intrapulmonary injection

Dose-related increases in the incidence of malignant lung tumours (mainly epidermoid and squamous-cell carcinomas and a few pleomorphic sarcomas) were found after injection of benzo[a]pyrene into the lung of rats ([Deutsch-Wenzel et al., 1983](#); [Iwagawa et al., 1989](#); [Wenzel-Hartung et al., 1990](#); [Horikawa et al., 1991](#)).

3.7 Intratracheal administration

Intratracheal administration of benzo[a]pyrene alone or mixed with particulates and suspended in saline with or without suspensions resulted in benign and malignant respiratory tumours in mice ([Heinrich et al., 1986a](#)), rats ([Pott et al., 1987](#); [Steinhoff et al., 1991](#)) and in numerous studies in hamsters ([IARC, 2010](#)). This treatment also induced forestomach tumours in hamsters ([Saffiotti et al., 1972](#); [Sellakumar et al., 1973](#); [Smith et al., 1975a, b](#); [Stenbäck & Rowland, 1979](#)). Larger benzo[a]pyrene particles were generally more effective than smaller ones.

Mice that lack the nucleotide excision-repair gene *XPA* (*XPA*^{-/-} mice) showed a stronger lung-tumour response after intratracheal instillation of benzo[a]pyrene than did their similarly treated *XPA*^{+/+} and *XPA*^{+/-} counterparts ([Ide et al., 2000](#)).

3.8 Buccal pouch application

Repeated application of benzo[a]pyrene to the buccal pouch mucosa of male hamsters resulted in a high incidence of forestomach papillomas ([Solt *et al.*, 1987](#)).

3.9 Subcutaneous tracheal grafts transplantation

In one study conducted in rats transplanted with subcutaneous rat tracheal grafts exposed to beeswax pellets containing various amounts of benzo[a]pyrene, a high incidence of squamous-cell carcinomas was reported ([Nettesheim *et al.*, 1977](#)).

3.10 Intramammary administration

In three studies in rats, benign and malignant mammary gland tumours developed after intramammary injection of benzo[a]pyrene ([Cavalieri *et al.*, 1988a, b, 1991](#)).

3.11 Intracolonic instillation

In three experiments in mice, intracolonic instillation of benzo[a]pyrene induced lymphomas and a variety of benign and malignant tumours in various organs including the forestomach ([Toth, 1980](#); [Anderson *et al.*, 1983](#)).

3.12 Intravaginal application

Intravaginal application of benzo[a]pyrene in mice produced invasive cervical carcinoma; no such tumours were seen in controls ([Näslund *et al.*, 1987](#)).

3.13 Intrafetal injection

In one study in male and female Swiss mice, intrafetal injection of benzo[a]pyrene produced lung adenomas ([Rossi *et al.*, 1983](#)).

4. Other Relevant Data

Benzo[a]pyrene is a carcinogen that induces tumours in many animal species. Some of the examples relevant for this review are: lung tumours in mice, rats, and hamsters; skin tumours in mice; liver tumours in mice; forestomach tumours in mice and hamsters; and mammary gland tumours in rats ([Osborne & Crosby, 1987](#); [IARC, 2010](#)). In humans, occupational exposures to benzo[a]pyrene-containing mixtures have been associated with a series of cancers: coke production: lung; coal gasification: lung, bladder; paving and roofing: lung; coal tar distillation: skin; soots: lung, oesophagus, haematolymphatic system, skin; aluminum smelting: lung, bladder; tobacco smoking: lung, lip, oral cavity, pharynx, oesophagus, larynx, bladder ([IARC, 1984, 1985, 1986, 2010](#)).

Studies on the mechanisms of action of benzo[a]pyrene have been reviewed ([Xue & Warshawsky, 2005](#); [IARC, 2010](#)).

4.1 Metabolism

Benzo[a]pyrene is metabolized by both phase-I and phase-II enzymes to form a series of arene oxides, dihydrodiols, phenols, and quinones and their polar conjugates with glutathione, sulfate, and glucuronide ([Osborne & Crosby, 1987](#)). Benzo[a]pyrene-7,8-diol is a key metabolite that is formed by the action of epoxide hydrolase on benzo[a]pyrene-7,8-epoxide. This dihydrodiol can be further metabolized by cytochrome P450s (CYPs) to a series of benzo[a]pyrene-7,8-diol-9,10-epoxides, which form one class of ultimate carcinogenic metabolites of benzo[a]pyrene.

Both CYPs and peroxidases (e.g. prostaglandin-H synthase) can oxidize benzo[*a*]pyrene. The major cytochrome P450s involved in the formation of diols and diolepoxides are CYP1A1, CYP1A2 and CYP1B1 ([Eling *et al.*, 1986](#); [Shimada, 2006](#)). Cytochrome P450s are inducible by benzo[*a*]pyrene and other PAHs through binding to the aryl hydrocarbon-receptor (AhR) nuclear complex, leading to changes in gene transcription of CYPs and phase-II enzymes. Mice lacking the AhR receptor are refractory to benzo[*a*]pyrene-induced tumorigenesis ([Shimizu *et al.*, 2000](#)). Both CYPs and peroxidases can form radical cations by one-electron oxidation. These cations comprise another class of ultimate carcinogenic metabolites ([Cavalieri & Rogan, 1995](#)). Some polymorphisms in human CYPs and phase-II enzymes (glutathione S-transferases, uridine 5'-diphosphate glucuronosyltransferases and sulfotransferases modulate susceptibility to cancer ([Shimada, 2006](#)). In another metabolic pathway, benzo[*a*]pyrene-7,8-dihydrodiol is oxidized to benzo[*a*]pyrene-7,8-quinone by enzymes of the aldo-keto reductase (AKR1) family. Among these, gene polymorphisms that influence susceptibility have been identified. NAD(P)H: quinone oxidoreductase-1 (NQO1) catalyses the reduction of benzo[*a*]pyrene quinones to hydroquinones, which may be re-oxidized and generate reactive oxygen species. Polymorphisms in this gene have also been described ([Penning & Drury, 2007](#); [IARC, 2010](#)).

The current understanding of mechanisms underlying benzo[*a*]pyrene-induced carcinogenesis in experimental animals is almost solely based on two complementary pathways: those of the diolepoxides and the radical cations. Each provides a different explanation for the effects observed in experimental animals in specific tissues.

4.2 Diolepoxide mechanism

The diolepoxide mechanism for benzo[*a*]pyrene features a sequence of metabolic transformations: benzo[*a*]pyrene → benzo[*a*]pyrene-7,8-oxide (by CYP1A1 and CYP1B1) → benzo[*a*]pyrene-7,8-diol (by epoxide hydrolase) → benzo[*a*]pyrene-7,8-diol-9,10-epoxides (by CYP1A1 and CYP1B1) ([Xue & Warshawsky, 2005](#)). Each class of metabolic intermediate has been shown to be genotoxic and carcinogenic ([Osborne & Crosby, 1987](#)). The stereochemistry of the metabolic transformation of benzo[*a*]pyrene to diols and diolepoxides is an important component of this mechanism of action. Due to the creation of chiral carbons during the metabolic conversions, many of the metabolic intermediates of benzo[*a*]pyrene have multiple stereochemical forms (enantiomeric and diastereomeric). As the metabolism proceeds the complexity of the stereo-chemical forms increases, eventually leading to four benzo[*a*]pyrene-7,8-diol-9,10-epoxides [(+)- and (-)-*anti*, (+)- and (-)-*syn*]. Diolepoxides react with DNA, mainly with the purines, deoxyguanosine and deoxyadenosine, and each diol-epoxide can form both *cis* and *trans* adducts thus giving a total of 16 possible benzo[*a*]pyrene-7,8-diol-9,10-epoxide DNA adducts. However, in most cases far fewer DNA adducts are actually observed. The most ubiquitous benzo[*a*]pyrene adduct detected in isolated mammalian DNA after metabolic conversion in metabolically competent mammalian cells in culture, or in mammals, is the *N*²-deoxyguanosine adduct, (+)-*N*²-10S-(7*R*,8*S*,9*R*-trihydroxy-7,8,9,10-tetrahydrobenzo[*a*]pyrene-yl)-2'-deoxyguanosine (BPDE-deoxyguanosine), derived from 7*R*,8*S*-dihydroxy-9*R*,10*R*-epoxy-7,8,9,10-tetrahydrobenzo[*a*]pyrene (*anti*-benzo[*a*]pyrene-7,8-diol-9,10-epoxide, or BPDE). This adduct was first fully identified after isolation from benzo[*a*]pyrene-treated human and bovine bronchial explants ([Jeffrey *et al.*, 1977](#)). This diolepoxide is considered to be an ultimate, DNA-reactive,

metabolite of benzo[a]pyrene ([Osborne & Crosby, 1987](#)). The *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide can form both stable and unstable (so-called 'depurinating') adducts with DNA, mediated by electrophilic carbonium ions ([Chakravarti et al., 2008](#)). *In vivo*, *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide produces stable adducts that were formed primarily with guanines in many species and organs ([IARC, 2010](#)).

Mice treated with benzo[a]pyrene had *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide-*N*²-deoxyguanosine adducts in their lung tissue, while the lung tumours induced by benzo[a]pyrene had G→T and G→A mutations in the *K_i-Ras* gene at codon 12 ([Mass et al., 1993](#)). In mice treated with benzo[a]pyrene the major stable DNA adduct in the epidermis was the *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-deoxyguanosine adduct ([Melendez-Colon et al., 1999](#)). Skin tumours from benzo[a]pyrene-treated mice or in preneoplastic skin from benzo[a]pyrene-treated mice had G→T mutations in codon 13 and A→T mutations in codon 61 of the *Ha-Ras* gene ([Chakravarti et al., 2008](#)).

Benzo[a]pyrene-induced skin tumours harboured G→T transversion mutations in the *Tp53* tumour-suppressor gene ([Ruggeri et al., 1993](#)). The *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adducts occurred at guanine positions in codons 157, 248, and 273 of the *TP53* gene in *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide-treated human HeLa cells. The same positions are the major mutational hotspots found in human lung cancers ([Denissenko et al., 1996](#)).

4.3 Radical-cation mechanism

The radical-cation mechanism for benzo[a]pyrene has been studied exclusively in connection with mouse-skin tumorigenesis ([Cavalieri & Rogan, 1995](#)). One-electron oxidation of benzo[a]pyrene by CYPs or peroxidases creates a radical cation localized on carbon 6, as a consequence of

the ionization potential and geometric configuration. In mouse skin, this radical cation gives rise to the formation of covalent adducts with guanine (at the C8 carbon and N7 nitrogen) and adenine (at the N7 nitrogen). These adducts are unstable and are thought to generate apurinic sites in mouse skin. However, only low levels of apurinic sites were measured in the epidermis of mice treated with benzo[a]pyrene ([Melendez-Colon et al., 1999](#)) and no studies to date have shown an increase in the number of apurinic sites in lung tissues treated with benzo[a]pyrene. In two *in vivo* studies, rats treated intraperitoneally with benzo[a]pyrene were shown to excrete 7-(benzo[a]pyrene-6-yl)-*N*7-guanine in faeces and urine, while the same adduct was detected in lung tissue of mice treated intraperitoneally with benzo[a]pyrene ([Rogan et al., 1990](#); [Banasiewicz et al., 2004](#)). Skin papillomas obtained from mice treated topically with benzo[a]pyrene showed mutations (at guanine and/or adenine) at codons 12, 13 and 61 in the *Ha-Ras* oncogene ([Wei et al., 1999](#)). Similar studies in preneoplastic skin from benzo[a]pyrene-treated mice showed *Ha-Ras* mutations at codons 13 and 61 ([Chakravarti et al., 2008](#)). The *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide can also form depurinating DNA adducts at guanine and adenine (at the N7 nitrogen). The distribution and chemical nature of the depurinating adducts (from both radical-cation and diolepoxide intermediates) in mouse skin and the distribution and chemical nature of the specific benzo[a]pyrene-induced mutations in mouse-skin papillomas have been reported ([Chakravarti et al., 2008](#)).

4.4 Other activation mechanisms of benzo[a]pyrene

4.4.1 Meso-region mechanism

The mechanism of meso-region biomethylation and benzylic oxidation features biomethylation of benzo[a]pyrene to 6-methylbenzo[a]

pyrene, with S-adenosylmethione as the carbon donor (Flesher *et al.*, 1982). This process has been shown to occur *in vitro*, and *in vivo* in rat liver (Stansbury *et al.*, 1994). 6-Methylbenzo[a]pyrene is further metabolized by CYPs to 6-hydroxymethylbenzo[a]pyrene (Flesher *et al.*, 1997) and then conjugated to sulfate by 3'-phosphoadenosine-5'-phosphosulfate sulfotransferase to 6-[(sulfooxy)methyl]-benzo[a]pyrene. This reactive sulfate ester forms DNA adducts *in vivo* (Stansbury *et al.*, 1994). These benzo[a]pyrene-DNA adducts have only been measured in rat liver (Surh *et al.*, 1989), which is not a target for benzo[a]pyrene-induced carcinogenesis. There is no evidence to date that this mechanism operates in lung.

4.4.2 Mechanism via formation of ortho-quinone/ reactive oxygen species

This mechanism features enzymatic oxidation of benzo[a]pyrene-7,8-diol to the *ortho*-quinone, benzo[a]pyrene-7,8-quinone, by aldo-keto reductases (Mangal *et al.*, 2009). Benzo[a]pyrene-7,8-quinone can react with DNA to yield both stable and depurinating DNA adducts *in vitro* (McCoull *et al.*, 1999; Balu *et al.*, 2006) and can also undergo repetitive redox cycling which generates reactive oxygen species that damage DNA (Penning *et al.*, 1999). In human A549 lung-tumour cells benzo[a]pyrene-7,8-quinone increased the formation of 8-oxo-deoxyguanosine and DNA strand-breaks (Park *et al.*, 2008; Mangal *et al.*, 2009). In a yeast reporter-assay, benzo[a]pyrene-7,8-quinone (in the presence of redox cycling) induced 8-oxo-deoxyguanosine formation and G→T transversions in the *Tp53* tumour-suppressor gene. The mutational spectra induced in the yeast reporter-assay closely matched those seen in DNA from human lung tumours (Shen *et al.*, 2006). Benzo[a]pyrene-7,8-quinone inhibited the activity of protein kinase C in MCF-7 cell lysates suggesting an ability to alter cell signalling (Yu *et al.*, 2002). Rats treated

with benzo[a]pyrene showed increased urinary concentrations of 8-oxo-deoxyguanosine, but lower levels in liver and lung tissues. This suggested that reactive oxygen species are generated during the CYP-dependent metabolism of benzo[a]pyrene, but induction of DNA-repair mechanisms may reduce these levels in target tissues (Briedé *et al.*, 2004). To date this mechanism has been studied only in *in-vitro* systems.

It is noted that formation of reactive oxygen species is not limited to the redox cycling of the *ortho*-quinone of benzo[a]pyrene (benzo[a]pyrene-7,8-quinone). There are several other sources of benzo[a]pyrene-induced reactive oxygen species. *In vivo*, both mice and rats metabolize benzo[a]pyrene to benzo[a]pyrene-1,6-quinone, benzo[a]pyrene-3,6-quinone and benzo[a]pyrene-6,12-quinone and these quinones enter into redox cycling and induce mutations (Osborne & Crosby, 1987; Joseph & Jaiswal, 1998). Many of the reactive intermediates of benzo[a]pyrene (oxides, diol-epoxides, radical cations) and quinone-generated reactive oxygen species can disrupt the balance of cellular oxidants and anti-oxidants by reducing the anti-oxidant levels thus leading to an imbalance and an excess of reactive oxygen species.

4.4.3 Aryl hydrocarbon-receptor mechanism

The AhR regulates the transcription of a series of genes including *Cyp1A1*, *Cyp1A2*, *Nqo1*, *Aldh3a1* (encoding aldehyde dehydrogenase 3A1), *UGT1a6* (uridine 5'-diphosphate-glucuronosyl transferase), and *Gsta1* (glutathione S-transferase A1). All these genes are activated by AhR-ligands, including benzo[a]pyrene, via the AhR-mediated aromatic hydrocarbon response element. The AhR plays a role in the response to oxidative stress in cell-cycle regulation and apoptosis. In addition, the CYP1A1/1A2-mediated metabolism generates oxidative stress (Nebert *et al.*, 2000). Mitochondrial hydrogen-peroxide production was induced by an AhR-ligand in

wild-type mice but not in *AhR*^{-/-} knockout mice (Senft *et al.*, 2002). These mice were shown to be refractory to benzo[a]pyrene-induced carcinogenicity (Shimizu *et al.*, 2000). Benzo[a]pyrene induced oxidative stress in mouse lung (Rajendran *et al.*, 2008).

4.4.4 Immunosuppression mechanism

Benzo[a]pyrene induces immunosuppression in adult mice by altering the cell-mediated responses (Wojdani & Alfred, 1984). Immune development in offspring is also altered following *in utero* exposure to benzo[a]pyrene (Urso & Gengoian, 1984). It is postulated that PAHs, including benzo[a]pyrene, act principally through their AhR-mediated CYP-derived metabolites (diolepoxides, quinones) to activate oxidative and electrophilic signalling pathways in lymphoid and nonlymphoid cells, including myeloid cells, epithelial cells, and other cell types. Furthermore, there is evidence that PAHs suppress immunity by p53-dependent pathways, by modulating signalling pathways in lymphocytes via non-genotoxic mechanisms, and by oxidative stress (Burchiel & Luster, 2001).

4.4.5 Epigenetic mechanisms

Benzo[a]pyrene and/or its metabolites have been shown to increase cell proliferation in several human cell lines, including terminally differentiated human bronchial squamous epithelial cells and in lung-cancer cells where increased expression of the *Cdc25B* gene (cell-division cycle 25B) was observed, along with reduced phosphorylation of Cdk1 (cyclin-dependent kinase 1) (Oguri *et al.*, 2003). Treatment with benzo[a]pyrene increased the number of human embryo lung-fibroblasts in the G1-S transition via the activation of c-Jun, through the p53-dependent PI-3K/Akt/ERK (phosphatidylinositol-3-kinase/protein kinase β /extracellular signal-regulated kinase) pathway (Jiao *et al.*, 2008).

Benzo[a]pyrene and/or its metabolites also affect apoptosis. Benzo[a]pyrene induced apoptosis in human MRC-5 lung fibroblasts via the JNK1/FasL (c-Jun N-terminal kinase 1/Fas Ligand) and JNK1/p53 signalling pathways (Chen *et al.*, 2005). Apoptosis induced by *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide in H460 human lung-cancer cells was associated with induction of Bak (BCL2-antagonist/killer) and with activation of caspase, but it was independent of Bcl-2 (Xiao *et al.*, 2007).

Altered DNA methylation has been reported to be associated with exposure to benzo[a]pyrene and/or its metabolites. After treatment of immortalized bronchial epithelial cells with *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide, the concentration of cytosine-DNA methyltransferase-1 was increased and was associated with hypermethylation of the promoters of 5–10 genes, including members of the cadherin gene-family (Damiani *et al.*, 2008).

4.5 Human exposure to PAH-rich mixtures

4.5.1 Biomarkers of exposure and effect

Molecular-epidemiological studies of cancer associated with occupational and environmental exposures to PAH have provided biomarkers that may be used to estimate internal exposure as well as to inform about molecular mechanisms that may be relevant to cancer causation, particularly in defining the early events in the carcinogenesis process. Biomarkers can be detected in the target organ, in surrogate tissues, or in tumours. These can be categorized into *biomarkers of exposure*, which are generally specific to the PAH of concern (e.g. DNA or protein adducts), *biomarkers of effect* (e.g. genotoxic and cytogenetic effects, 8-oxo-deoxyguanosine, sister chromatid exchange (SCE), micronuclei, chromosomal aberrations, mutations in oncogenes, tumour-suppressor genes, or indicator genes),

and *biomarkers of susceptibility* (DNA-repair enzymes, e.g. XPA, XPC – *xeroderma pigmentosum* complementation groups A and C), bioactivation enzymes (e.g. CYPs), detoxification enzymes (e.g. GSTs), and mutagenic metabolites in urine ([Kalina et al., 1998](#); [Pilger et al., 2000](#); [Simioli et al., 2004](#); [Raimondi et al., 2005](#); [Vineis & Husgafvel-Pursiainen, 2005](#); [Matullo et al., 2006](#); [Farmer & Singh, 2008](#); [Gyorffy et al., 2008](#)). Although biomarkers of effect and susceptibility are generally not unique to any specific PAH exposure, several these biomarkers may provide insight into the mechanism of carcinogenesis induced in humans by PAHs or PAH-rich exposures.

4.5.2 Exposure to benzo[a]pyrene and relationship with specific biomarkers

Biomarkers of exposure to complex mixtures that contain benzo[a]pyrene have been studied in populations exposed in industrial settings: coke production, coal-tar distillation, the aluminium industry, roofing and paving with coal-tar pitch, coal gasification, chimney sweeping, and iron and steel founding. Most if not all of these biomarkers are genotoxic markers. Populations of patients who undergo coal-tar therapy and groups exposed to combustion emissions, and tobacco smokers have also been evaluated. Studies on biomarkers of exposure are dominated by those focusing on the *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adduct, the most commonly studied PAH-DNA adduct because of the availability of specific analytical methods and standards ([Gyorffy et al., 2008](#)). In one study the depurinating adducts resulting from radical-cation formation, *viz.* 7-(benzo[a]pyrene-6-yl)guanine and 7-(benzo[a]pyrene-6-yl)adenine were found in the urine of women exposed to coal smoke ([Casale et al., 2001](#)). Concomitantly, several biomarkers of effect have also been evaluated in these studies: chromosomal aberrations, sister chromatid exchange ([Kalina et al., 1998](#)),

DNA damage (measured by the comet assay) and 8-oxo-deoxyguanosine formation ([Marczynski et al., 2002](#)). It is important to note that these genotoxic effects observed in humans in relation to exposure to benzo[a]pyrene-containing mixtures have also been observed in experimental studies where benzo[a]pyrene or *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide has been shown to induce sister chromatid exchange ([Pal et al., 1980](#); [Brauze et al., 1997](#)), chromosomal aberrations, micronuclei ([Kliesch et al., 1982](#)), DNA damage ([Nesnow et al., 2002](#)), and 8-oxo-deoxyguanosine ([Thaiparambil et al., 2007](#)). Tobacco smoke, dietary habits and indoor ambient air are also important sources of exposure to benzo[a]pyrene, which has been implicated as one of the components of tobacco smoke related to the induction of lung cancer in smokers ([Watanabe et al., 2009](#)). In a large study of 585 smokers and nonsmokers, smoking and diet were highly correlated with *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adduct levels ([Pavanello et al., 2006](#)). Several studies have demonstrated moderately increased levels of 8-oxo-deoxyguanosine from lungs, sperm, and leukocytes of smokers. Increased urinary excretion of 8-oxo-deoxyguanosine has also been reported ([Hecht, 1999](#)). In rats exposed to benzo[a]pyrene via oral, intratracheal and dermal routes, *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adducts were formed in white blood cells independently of the exposure route and their numbers correlated with those found in lung DNA, suggesting that *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adduct levels in white blood cells may be used as a surrogate for pulmonary *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adducts ([Godschalk et al., 2000](#)).

4.5.3 Relationship of biomarkers to human cancer

Mutations in *TP53* are common in lung cancers from smokers and less common in nonsmokers. These mutations are G→T transversions with hotspots in codons 157, 248 and 273 ([Hainaut & Pfeifer, 2001](#); [Pfeifer et al., 2002](#)) and they are associated with *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adducts. The active metabolite *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide causes a unique spectrum of *TP53* mutations distinct from those found in cancers that are not associated with smoking ([Campling & el-Deiry, 2003](#)). Similar G→T mutations have been reported in lung tumours from nonsmoking Chinese women whose tumours were associated with exposure to PAHs from smoke generated by burning smoky coal in unventilated homes. The mutations were clustered at the CpG rich codons 153–158 of the *TP53* gene, and at codons 249 and 273. The mutation spectrum was fully consistent with exposure to PAHs ([DeMarini et al., 2001](#)).

4.6 Synthesis

Benzo[a]pyrene is metabolically activated to a series of reactive intermediates by CYP450 and related enzymes under control of the aryl-hydrocarbon receptor. There is strong evidence that the benzo[a]pyrene diolepoxide mechanism operates in mouse-lung tumorigenesis, while there is also strong evidence that both the radical-cation and the diolepoxide mechanisms are involved in mouse-skin carcinogenesis. The meso-region mechanism has been studied only in rat liver, while the mechanism that involves the formation of *ortho*-quinone/reactive oxygen species has only been studied *in vitro*, although reactive oxygen species can be formed *in vivo* by other benzo[a]pyrene-mediated mechanisms. All these pathways reflect genotoxic mechanisms, as they involve alterations to DNA. Benzo[a]pyrene is pleiotropic and has the ability to affect many

cell- and organ-based systems. Therefore, there are probably many modes of carcinogenic action operating to different extents *in vivo*. These include mechanisms that involve AhR, oxidative stress, immunotoxicity and epigenetic events.

Based on the best available, consistent and strong experimental and human mechanistic evidence it is concluded that benzo[a]pyrene contributes to the genotoxic and carcinogenic effects resulting from occupational exposure to complex PAH mixtures that contain benzo[a]pyrene. The most commonly encountered – and most widely studied – mechanistically relevant DNA lesion is the *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adduct. The formation of this adduct is consistent with *anti*-benzo[a]pyrene-7,8-diol-9,10-epoxide-associated genotoxic effects in surrogate tissues and with the mutation pattern in the *TP53* gene in lung tumours from humans exposed to PAH mixtures that contain benzo[a]pyrene. The fact that those PAH mixtures and benzo[a]pyrene itself induce genotoxic effects like sister chromatid exchange, chromosomal aberrations, micronuclei, DNA damage (comet assay) and 8-oxo-deoxyguanosine, supports the notion that benzo[a]pyrene contributes to human cancer.

5. Evaluation

There is *sufficient evidence* for the carcinogenicity of benzo[a]pyrene in experimental animals.

[No epidemiological data on benzo[a]pyrene alone were available to the Working Group.]

The genotoxic mechanism of action of benzo[a]pyrene involves metabolism to highly reactive species that form covalent adducts to DNA. These *anti*-benzo[a]pyrene-7,8-diol-9,10-oxide-DNA adducts induce mutations in the *K-RAS* oncogene and the *TP53* tumour-suppressor gene in human lung tumours, and

in corresponding genes in mouse-lung tumours. Exposure to benzo[a]pyrene and benzo[a]pyrene-containing complex mixtures also induce other genotoxic effects, including sister chromatid exchange, micronuclei, DNA damage and 8-oxo-deoxyguanosine, all of which can contribute to the carcinogenic effects of benzo[a]pyrene and benzo[a]pyrene-containing complex mixtures in exposed humans.

Benzo[a]pyrene is *carcinogenic to humans* (Group 1).

In making the overall evaluation, the Working Group took the following into consideration:

The strong and extensive experimental evidence for the carcinogenicity of benzo[a]pyrene in many animal species, supported by the consistent and coherent mechanistic evidence from experimental and human studies provide biological plausibility to support the overall classification of benzo[a]pyrene as a human carcinogen (Group 1).

References

- Albert RE, Miller ML, Cody T *et al.* (1991). Benzo[a]pyrene-induced skin damage and tumor promotion in the mouse. *Carcinogenesis*, 12: 1273–1280. doi:10.1093/carcin/12.7.1273 PMID:2070493
- Anderson LM, Priest LJ, Deschner EE, Budinger JM (1983). Carcinogenic effects of intracolonic benzo[a]pyrene in β -naphthoflavone-induced mice. *Cancer Letters*, 20: 117–123. doi:10.1016/0304-3835(83)90039-3 PMID:6321017
- Andrews J, Halliday GM, Muller HK (1991). A role for prostaglandins in the suppression of cutaneous cellular immunity and tumour development in benzo[a]pyrene-but not dimethylbenz(a)anthracene-treated mice. *Clin Exp Immunol*, 85: 9–13. PMID:1906386
- Badary OA, Al-Shabanah OA, Nagi MN *et al.* (1999). Inhibition of benzo[a]pyrene-induced forestomach carcinogenesis in mice by thymoquinone. *European Journal of Cancer Prevention*, 8: 435–440. doi:10.1097/00008469-199910000-00009 PMID:10548399
- Balansky R, Ganchev G, Ilcheva M *et al.* (2007). Potent carcinogenicity of cigarette smoke in mice exposed early in life. *Carcinogenesis*, 28: 2236–2243. doi:10.1093/carcin/bgm122 PMID:17522065
- Balu N, Padgett WT, Nelson GB *et al.* (2006). Benzo[a]pyrene-7,8-quinone-3'-mononucleotide adduct standards for 32P postlabeling analyses: detection of benzo[a]pyrene-7,8-quinone-calf thymus DNA adducts. *Anal Biochem*, 355: 213–223. doi:10.1016/j.ab.2006.05.023 PMID:16797471
- Banasiewicz M, Nelson G, Swank A *et al.* (2004). Identification and quantitation of benzo[a]pyrene-derived DNA adducts formed at low adduction level in mice lung tissue. *Anal Biochem*, 334: 390–400. doi:10.1016/j.ab.2004.08.006 PMID:15494147
- Brauze D, Wielgosz SM, Pawlak AL, Baer-Dubowska W (1997). Effect of the route of benzo[a]pyrene administration on sister chromatid exchange and DNA binding in bone marrow of mice differing with respect to cytochrome P450 1A1 induction. *Toxicol Lett*, 91: 211–217. doi:10.1016/S0378-4274(97)00024-6 PMID:9217241
- Briedé JJ, Godschalk RW, Emans MT *et al.* (2004). In vitro and in vivo studies on oxygen free radical and DNA adduct formation in rat lung and liver during benzo[a]pyrene metabolism. *Free Radic Res*, 38: 995–1002. doi:10.1080/10715760400000976 PMID:15621718
- Burchiel SW & Luster MI (2001). Signalling by environmental polycyclic aromatic hydrocarbons in human lymphocytes. *Clin Immunol*, 98: 2–10. doi:10.1006/clim.2000.4934 PMID:11141320
- Busby WF Jr, Stevens EK, Martin CN *et al.* (1989). Comparative lung tumorigenicity of parent and mononitro-polynuclear aromatic hydrocarbons in the BLU:Ha newborn mouse assay. *Toxicology and Applied Pharmacology*, 99: 555–563. doi:10.1016/0041-008X(89)90162-2 PMID:2749740
- Campling BG & el-Deiry WS (2003). Clinical implications of p53 mutations in lung cancer. *Methods Mol Med*, 75: 53–77. PMID:12407735
- Casale GP, Singhal M, Bhattacharya S *et al.* (2001). Detection and quantification of depurinated benzo[a]pyrene-adducted DNA bases in the urine of cigarette smokers and women exposed to household coal smoke. *Chem Res Toxicol*, 14: 192–201. doi:10.1021/tx000012y PMID:11258968
- Cavalieri E, Mailander P, Pelfrene A (1977). Carcinogenic activity of anthanthrene on mouse skin. *Zeitschrift für Krebsforschung*, 89: 113–118. PMID:143140 doi:10.1007/BF00308512
- Cavalieri E, Rogan E, Cremonesi P *et al.* (1988a). Tumorigenicity of 6-halogenated derivatives of benzo[a]pyrene in mouse skin and rat mammary gland. *Journal of Cancer Research and Clinical Oncology*, 114: 10–15. doi:10.1007/BF00390479 PMID:3350835
- Cavalieri E, Rogan E, Sinha D (1988b). Carcinogenicity of aromatic hydrocarbons directly applied to rat mammary gland. *J. Cancer clin. Oncol*, 114: 3–9. PMID:3350839.
- Cavalieri EL, Higginbotham S, RamaKrishna NVS *et al.* (1991). Comparative dose-response

- tumorigenicity studies of dibenzo[a,l]pyrene versus 7,12-dimethylbenz[a]anthracene, benzo[a]pyrene and two dibenzo[a,l]pyrene dihydrols in mouse skin and rat mammary gland. *Carcinogenesis*, 12: 1939–1944. doi:10.1093/carcin/12.10.1939 PMID:1934274
- Cavalieri EL & Rogan EG (1995). Central role of radical cations in metabolic activation of polycyclic aromatic hydrocarbons. *Xenobiotica*, 25: 677–688. doi:10.3109/00498259509061885 PMID:7483666
- Chakravarti D, Venugopal D, Mailander PC *et al.* (2008). The role of polycyclic aromatic hydrocarbon-DNA adducts in inducing mutations in mouse skin. *Mutat Res*, 649: 161–178. PMID:17931959
- Chen JH, Chou FP, Lin HH, Wang CJ (2005). Gaseous nitrogen oxide repressed benzo[a]pyrene-induced human lung fibroblast cell apoptosis via inhibiting JNK1 signals. *Arch Toxicol*, 79: 694–704. doi:10.1007/s00204-005-0001-0 PMID:16041517
- Culp SJ, Gaylor DW, Sheldon WG *et al.* (1998). A comparison of the tumors induced by coal tar and benzo[a]pyrene in a 2-year bioassay. *Carcinogenesis*, 19: 117–124. doi:10.1093/carcin/19.1.117 PMID:9472702
- Damiani LA, Yingling CM, Leng S *et al.* (2008). Carcinogen-induced gene promoter hypermethylation is mediated by DNMT1 and causal for transformation of immortalized bronchial epithelial cells. *Cancer Res*, 68: 9005–9014. doi:10.1158/0008-5472.CAN-08-1276 PMID:18974146
- de Vries A, van Oostrom CTM, Dortant PM *et al.* (1997). Spontaneous liver tumors and benzo[a]pyrene-induced lymphomas in XPA-deficient mice. *Molecular Carcinogenesis*, 19: 46–53. doi:10.1002/(SICI)1098-2744(199705)19:1<46::AID-MC7>3.0.CO;2-L PMID:9180928
- DeMarini DM, Landi S, Tian D *et al.* (2001). Lung tumor KRAS and TP53 mutations in nonsmokers reflect exposure to PAH-rich coal combustion emissions. *Cancer Res*, 61: 6679–6681. PMID:11559534
- Denissenko MF, Pao A, Tang M, Pfeifer GP (1996). Preferential formation of benzo[a]pyrene adducts at lung cancer mutational hotspots in P53. *Science*, 274: 430–432. doi:10.1126/science.274.5286.430 PMID:8832894
- Deutsch-Wenzel RP, Brune H, Grimmer G *et al.* (1983). Experimental studies in rat lungs on the carcinogenicity and dose-response relationships of eight frequently occurring environmental polycyclic aromatic hydrocarbons. *J Natl Cancer Inst*, 71: 539–544. PMID:6577228
- el-Bayoumy K, Chae Y-H, Upadhyaya P *et al.* (1995). Comparative tumorigenicity of benzo[a]pyrene, 1-nitropyrene and 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine administered by gavage to female CD rats. *Carcinogenesis*, 16: 431–434. doi:10.1093/carcin/16.2.431 PMID:7859378
- Eling T, Curtis J, Battista J, Marnett LJ (1986). Oxidation of (+)-7,8-dihydroxy-7,8-dihydrobenzo[a]pyrene by mouse keratinocytes: evidence for peroxy radical- and monooxygenase-dependent metabolism. *Carcinogenesis*, 7: 1957–1963. doi:10.1093/carcin/7.12.1957 PMID:2430728
- Estensen RD, Jordan MM, Wiedmann TS *et al.* (2004). Effect of chemopreventive agents on separate stages of progression of benzo[alpha]pyrene induced lung tumors in A/J mice. *Carcinogenesis*, 25: 197–201. doi:10.1093/carcin/bgg196 PMID:14578161
- Estensen RD & Wattenberg LW (1993). Studies of chemopreventive effects of myo-inositol on benzo[a]pyrene-induced neoplasia of the lung and forestomach of female A/J mice. *Carcinogenesis*, 14: 1975–1977. doi:10.1093/carcin/14.9.1975 PMID:8403228
- Farmer PB & Singh R (2008). Use of DNA adducts to identify human health risk from exposure to hazardous environmental pollutants: the increasing role of mass spectrometry in assessing biologically effective doses of genotoxic carcinogens. *Mutat Res*, 659: 68–76. doi:10.1016/j.mrrev.2008.03.006 PMID:18468947
- Feron VJ (1972). Respiratory tract tumors in hamsters after intratracheal instillations of benzo(a)pyrene alone and with furfural. *Cancer Res*, 32: 28–36. PMID:5007686
- Feron VJ, de Jong D, Emmelot P (1973). Letter: Dose-response correlation for the induction of respiratory-tract tumours in Syrian golden hamsters by intratracheal instillations of benzo(a)pyrene. *Eur J Cancer*, 9: 387–390. doi:10.1016/0014-2964(73)90057-1 PMID:4746737
- Feron VJ & Kruijsse A (1978). Effects of exposure to furfural vapour in hamsters simultaneously treated with benzo[alpha]pyrene or diethylnitrosamine. *Toxicology*, 11: 127–144. doi:10.1016/S0300-483X(78)90889-2 PMID:715798
- Feron VJ, van den Heuvel PD, Koeter HB, Beems RB (1980). Significance of particle size of benzo(a)pyrene for the induction of respiratory tract tumours in hamsters. *Int J Cancer*, 25: 301–307. doi:10.1002/ijc.2910250220 PMID:7390654
- Flesher JW, Horn J, Lehner AF (1997). 6-sulfooxymethylbenzo[a]pyrene is an ultimate electrophilic and carcinogenic form of the intermediary metabolite 6-hydroxymethylbenzo[a]pyrene. *Biochem Biophys Res Commun*, 234: 554–558. doi:10.1006/bbrc.1997.6679 PMID:9175750
- Flesher JW, Stansbury KH, Sydnor KL (1982). S-Adenosyl-L-methionine is a carbon donor in the conversion of benzo[alpha]pyrene to 6-hydroxymethylbenzo[alpha]pyrene by rat liver S-9. *Cancer Lett*, 16: 91–94. doi:10.1016/0304-3835(82)90095-7 PMID:6288234
- Godleski JJ, Melnicoff MJ, Sadri S, Garbeil P (1984). Effects of inhaled ammonium sulfate on benzo[a]pyrene carcinogenesis. *J Toxicol Environ Health*, 14: 225–238. doi:10.1080/15287398409530575 PMID:6502734
- Godschalk RW, Moonen EJ, Schilderman PA *et al.* (2000). Exposure-route-dependent DNA adduct formation by

- polycyclic aromatic hydrocarbons. *Carcinogenesis*, 21: 87–92. doi:10.1093/carcin/21.1.87 PMID:10607738
- Gyorfy E, Anna L, Kovács K *et al.* (2008). Correlation between biomarkers of human exposure to genotoxins with focus on carcinogen-DNA adducts. *Mutagenesis*, 23: 1–18. doi:10.1093/mutage/gem043 PMID:17989146
- Habs M, Jahn SAA, Schmähl D (1984). Carcinogenic activity of condensate from colocynth seeds (*Citrullus colocynthis*) after chronic epicutaneous administration to mice. *J Cancer Res Clin Oncol*, 108: 154–156. doi:10.1007/BF00390988 PMID:6746706
- Habs M, Schmähl D, Misfeld J (1980). Local carcinogenicity of some environmentally relevant polycyclic aromatic hydrocarbons after lifelong topical application to mouse skin. *Arch Geschwulstforsch*, 50: 266–274. PMID:7436704
- Hainaut P & Pfeifer GP (2001). Patterns of p53 G→T transversions in lung cancers reflect the primary mutagenic signature of DNA-damage by tobacco smoke. *Carcinogenesis*, 22: 367–374. doi:10.1093/carcin/22.3.367 PMID:11238174
- Hakura A, Tsutsui Y, Sonoda J *et al.* (1998). Comparison between in vivo mutagenicity and carcinogenicity in multiple organs by benzo[a]pyrene in the lacZ transgenic mouse (Muta Mouse). *Mutat Res*, 398: 123–130. PMID:9626972
- Hecht SS (1999). Tobacco smoke carcinogens and lung cancer. *J Natl Cancer Inst*, 91: 1194–1210. doi:10.1093/jnci/91.14.1194 PMID:10413421
- Heinrich U, Pott F, Mohr U *et al.* (1986a). Lung tumours in rats and mice after inhalation of PAH-rich emissions. *Exp Pathol*, 29: 29–34. PMID:3699126
- Henry MC, Port CD, Bates RR, Kaufman DG (1973). Respiratory tract tumors in hamsters induced by benzo(a)pyrene. *Cancer Res*, 33: 1585–1592. PMID:4721222
- Homburger F, Hsueh S-S, Kerr CS, Russfield AB (1972). Inherited susceptibility of inbred strains of Syrian hamsters to induction of subcutaneous sarcomas and mammary and gastrointestinal carcinomas by subcutaneous and gastric administration of polynuclear hydrocarbons. *Cancer Res*, 32: 360–366. PMID:5058191
- Hoogervorst EM, de Vries A, Beems RB *et al.* (2003). Combined oral benzo[a]pyrene and inhalatory ozone exposure have no effect on lung tumor development in DNA repair-deficient Xpa mice. *Carcinogenesis*, 24: 613–619. doi:10.1093/carcin/24.3.613 PMID:12663525
- Horikawa K, Sera N, Otofujii T *et al.* (1991). Pulmonary carcinogenicity of 3,9- and 3,7-dinitrofluoranthene, 3-nitrofluoranthene and benzo[a]pyrene in F344 rats. *Carcinogenesis*, 12: 1003–1007. doi:10.1093/carcin/12.6.1003 PMID:2044179
- IARC (1973). Certain polycyclic aromatic hydrocarbons and heterocyclic compounds. *IARC Monogr Eval Carcinog Risk Chem Man*, 3: 1–271.
- IARC (1983). Polynuclear aromatic compounds, Part 1, chemical, environmental and experimental data. *IARC Monogr Eval Carcinog Risk Chem Hum*, 32: 1–453. PMID:6586639
- IARC (1984). Polynuclear aromatic compounds, Part 3, industrial exposures in aluminium production, coal gasification, coke production, and iron and steel founding. *IARC Monogr Eval Carcinog Risk Chem Hum*, 34: 1–219.
- IARC (1985). Polynuclear aromatic compounds, Part 4, bitumens, coal-tars and derived products, shale-oils and soots. *IARC Monogr Eval Carcinog Risk Chem Hum*, 35: 1–247. PMID:2991123
- IARC (1986). Tobacco smoking. *IARC Monogr Eval Carcinog Risk Chem Hum*, 38: 1–421.
- IARC (2010). Some non-heterocyclic polycyclic aromatic hydrocarbons and some related exposures. *IARC Monogr Eval Carcinog Risks Hum*, 92: 1–853. PMID:21141735 PMID:18756632
- Ide F, Iida N, Nakatsuru Y *et al.* (2000). Mice deficient in the nucleotide excision repair gene XPA have elevated sensitivity to benzo[a]pyrene induction of lung tumors. *Carcinogenesis*, 21: 1263–1265. doi:10.1093/carcin/21.6.1263 PMID:10837020
- Iwagawa M, Maeda T, Izumi K *et al.* (1989). Comparative dose-response study on the pulmonary carcinogenicity of 1,6-dinitropyrene and benzo[a]pyrene in F344 rats. *Carcinogenesis*, 10: 1285–1290. doi:10.1093/carcin/10.7.1285 PMID:2736719
- Jeffrey AM, Weinstein IB, Jennette KW *et al.* (1977). Structures of benzo(a)pyrene-nucleic acid adducts formed in human and bovine bronchial explants. *Nature*, 269: 348–350. doi:10.1038/269348a0 PMID:904688
- Jiao S, Liu B, Gao A *et al.* (2008). Benzo(a)pyrene-caused increased G1-S transition requires the activation of c-Jun through p53-dependent PI-3K/Akt/ERK pathway in human embryo lung fibroblasts. *Toxicol Lett*, 178: 167–175. doi:10.1016/j.toxlet.2008.03.012 PMID:18448277
- Joseph P & Jaiswal AK (1998). NAD(P)H:quinone oxidoreductase 1 reduces the mutagenicity of DNA caused by NADPH:P450 reductase-activated metabolites of benzo(a)pyrene quinones. *Br J Cancer*, 77: 709–719. PMID:9514048
- Kalina I, Brezáni P, Gajdosová D *et al.* (1998). Cytogenetic monitoring in coke oven workers. *Mutat Res*, 417: 9–17. PMID:9729241
- Ketkar M, Green U, Schneider P, Mohr U (1979). Investigations on the carcinogenic burden by air pollution in man. Intratracheal instillation studies with benzo[a]pyrene in a mixture of Tris buffer and saline in Syrian golden hamsters. *Cancer Lett*, 6: 279–284. doi:10.1016/S0304-3835(79)80046-4 PMID:436122
- Ketkar M, Reznik G, Misfeld J, Mohr U (1977). Investigations on the carcinogenic burden by air pollution in man. The effect of a single dose of benzo(a)

- pyrene in Syrian golden hamsters. *Cancer Lett*, 3: 231–235. doi:10.1016/S0304-3835(77)95980-8
- Ketkar M, Reznik G, Schneider P, Mohr U (1978). Investigations on the carcinogenic burden by air pollution in man. Intratracheal instillation studies with benzo(a)pyrene in bovine serum albumin in Syrian hamsters. *Cancer Lett*, 4: 235–239. doi:10.1016/S0304-3835(78)94787-0 PMID:647664
- Kliesch U, Roupova I, Adler ID (1982). Induction of chromosome damage in mouse bone marrow by benzo[a]pyrene. *Mutat Res*, 102: 265–273. doi:10.1016/0165-1218(82)90136-7 PMID:7144782
- Kobayashi N (1975). Production of respiratory tract tumors in hamsters by benzo(a)pyrene. *Gann*, 66: 311–315. PMID:1181231
- Kouri RE, Wood AW, Levin W *et al.* (1980). Carcinogenicity of benzo[a]pyrene and thirteen of its derivatives in C3H/FCum mice. *J Natl Cancer Inst*, 64: 617–623. PMID:6766516
- Kroese ED, Dortant PM, van Steeg H *et al.* (1997). Use of E μ -PIM-1 transgenic mice short-term in vivo carcinogenicity testing: lymphoma induction by benzo[a]pyrene, but not by TPA. *Carcinogenesis*, 18: 975–980. doi:10.1093/carcin/18.5.975 PMID:9163683
- Kruysse A & Feron VJ (1976). Repeated exposure to cyclopentenone vapour: long-term study in Syrian golden hamsters *Zentralbl Bakteriol*, [Orig.B]163: 5–6448–457. . PMID:1020535.
- Lavoie EJ, Braley J, Rice JE, Rivenson A (1987). Tumorigenic activity of non-alternant polynuclear aromatic hydrocarbons in newborn mice. *Cancer Letters*, 34: 15–20. doi:10.1016/0304-3835(87)90068-1 PMID:3802065
- Levin W, Wood AW, Wislocki PG *et al.* (1977). Carcinogenicity of benzo ring derivatives of benzo[a]pyrene on mouse skin. *Cancer Research*, 37: 3357–3361. PMID:884679.
- Mangal D, Vudathala D, Park JH *et al.* (2009). Analysis of 7,8-dihydro-8-oxo-2'-deoxyguanosine in cellular DNA during oxidative stress. *Chem Res Toxicol*, 22: 788–797. doi:10.1021/tx800343c PMID:19309085
- Marczynski B, Rihs HP, Rossbach B *et al.* (2002). Analysis of 8-oxo-7,8-dihydro-2'-deoxyguanosine and DNA strand breaks in white blood cells of occupationally exposed workers: comparison with ambient monitoring, urinary metabolites and enzyme polymorphisms. *Carcinogenesis*, 23: 273–281. doi:10.1093/carcin/23.2.273 PMID:11872632
- Mass MJ, Jeffers AJ, Ross JA *et al.* (1993). Ki-ras oncogene mutations in tumors and DNA adducts formed by benz[j]aceanthrylene and benzo[a]pyrene in the lungs of strain A/J mice. *Molecular Carcinogenesis*, 8: 186–192. doi:10.1002/mc.2940080309 PMID:8216737
- Matullo G, Dunning AM, Guarrera S *et al.* (2006). DNA repair polymorphisms and cancer risk in non-smokers in a cohort study. *Carcinogenesis*, 27: 997–1007. doi:10.1093/carcin/bgi280 PMID:16308313
- McCoull KD, Rindgen D, Blair IA, Penning TM (1999). Synthesis and characterization of polycyclic aromatic hydrocarbon o-quinone depurinating N7-guanine adducts. *Chem Res Toxicol*, 12: 237–246. doi:10.1021/tx980182z PMID:10077486
- Melendez-Colon VJ, Luch A, Seidel A, Baird WM (1999). Cancer initiation by polycyclic aromatic hydrocarbons results from formation of stable DNA adducts rather than apurinic sites. *Carcinogenesis*, 20: 1885–1891. doi:10.1093/carcin/20.10.1885 PMID:10506100
- Näslund I, Rubio CA, Auer GU (1987). Nuclear DNA changes during pathogenesis of squamous cell carcinoma of the cervix in 3,4-benzopyrene-treated mice. *Analytical and Quantitative Cytology*, 9: 411–418. PMID: 3675800.
- Nebert DW, Roe AL, Dieter MZ *et al.* (2000). Role of the aromatic hydrocarbon receptor and [Ah] gene battery in the oxidative stress response, cell cycle control, and apoptosis. *Biochem Pharmacol*, 59: 65–85. doi:10.1016/S0006-2952(99)00310-X PMID:10605936
- Nesnow S, Davis C, Nelson GB *et al.* (2002). Comparison of the genotoxic activities of the K-region dihydrodiol of benzo[a]pyrene with benzo[a]pyrene in mammalian cells: morphological cell transformation; DNA damage; and stable covalent DNA adducts. *Mutat Res*, 521: 91–102. PMID:12438007
- Nesnow S, Ross JA, Stoner GD, Mass MJ (1995). Mechanistic linkage between DNA adducts, mutations in oncogenes and tumorigenesis of carcinogenic environmental polycyclic aromatic hydrocarbons in strain A/J mice. *Toxicology*, 105: 403–413. doi:10.1016/0300-483X(95)03238-B PMID:8571376
- Nettesheim P, Griesemer RA, Martin DH, Caton JE Jr (1977). Induction of preneoplastic and neoplastic lesions in grafted rat tracheas continuously exposed to benzo[a]pyrene. *Cancer Research*, 37: 1271–1278. PMID: 856459.
- Oguri T, Singh SV, Nemoto K, Lazo JS (2003). The carcinogen (7R,8S)-dihydroxy-(9S,10R)-epoxy-7,8,9,10-tetrahydrobenzo[a]pyrene induces Cdc25B expression in human bronchial and lung cancer cells. *Cancer Res*, 63: 771–775. PMID:12591724
- Osborne MR, Crosby NT (1987). *Benzo[a]pyrenes*. Cambridge Monographs on Cancer Research. Cambridge, England: Cambridge University Press
- Pal K, Grover PL, Sims P (1980). The induction of sister-chromatid exchanges in Chinese hamster ovary cells by some epoxides and phenolic derivatives of benzo[a]pyrene. *Mutat Res*, 78: 193–199. doi:10.1016/0165-1218(80)90098-1 PMID:7393246
- Park JH, Mangal D, Tacka KA *et al.* (2008). Evidence for the aldo-keto reductase pathway of polycyclic aromatic trans-dihydrodiol activation in human lung A549 cells. *Proc Natl Acad Sci U S A*, 105: 6846–6851. doi:10.1073/pnas.0802776105 PMID:18474869

- Pavanello S, Pulliero A, Saia BO, Clonfero E (2006). Determinants of anti-benzo[a]pyrene diol epoxide-DNA adduct formation in lymphomonocytes of the general population. *Mutat Res*, 611: 54–63. PMID:16978913
- Penning TM, Burczynski ME, Hung CF *et al.* (1999). Dihydrodiol dehydrogenases and polycyclic aromatic hydrocarbon activation: generation of reactive and redox active o-quinones. *Chem Res Toxicol*, 12: 1–18. doi:10.1021/tx980143n PMID:9894013
- Penning TM & Drury JE (2007). Human aldo-keto reductases: Function, gene regulation, and single nucleotide polymorphisms. *Arch Biochem Biophys*, 464: 241–250. doi:10.1016/j.abb.2007.04.024 PMID:17537398
- Pfeifer GP, Denissenko MF, Olivier M *et al.* (2002). Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. *Oncogene*, 21: 7435–7451. doi:10.1038/sj.onc.1205803 PMID:12379884
- Pilger A, Germadnik D, Schaffer A *et al.* (2000). 8-Hydroxydeoxyguanosine in leukocyte DNA and urine of quartz-exposed workers and patients with silicosis. *Int Arch Occup Environ Health*, 73: 305–310. doi:10.1007/s004200000117 PMID:10963413
- Pott F, Brockhaus A, Huth F (1973a) [Tests on the production of tumours in animal experiments with polycyclic aromatic hydrocarbons.] [in German]. *Zbl. Bakt. Hyg. Abt. Orig. B*, 157: 34–43.
- Pott F, Tomingas R, Reiffer FJ (1973b). Experimental studies on the carcinogenicity and the retention of benzo[a]pyrene in application region after intratracheal and subcutaneous injection. *Zbl. Bakt. Hyg. I. Abt. Orig. B*, 158: 97–108.
- Pott F, Ziem U, Reiffer F-J *et al.* (1987). Carcinogenicity studies on fibres, metal compounds, and some other dusts in rats. *Exp Pathol*, 32: 129–152. PMID:3436395
- Raimondi S, Boffetta P, Anttila S *et al.* (2005). Metabolic gene polymorphisms and lung cancer risk in non-smokers. An update of the GSEC study. *Mutat Res*, 592: 45–57. PMID:16009381
- Rajendran P, Ekambaram G, Sakthisekaran D (2008). Cytoprotective effect of mangiferin on benzo(a) pyrene-induced lung carcinogenesis in swiss albino mice. *Basic Clin Pharmacol Toxicol*, 103: 137–142. doi:10.1111/j.1742-7843.2008.00254.x PMID:18816296
- Reynders JB, Immel HR, Scherrenberg PM *et al.* (1985). Respiratory tract tumors in hamsters after severe focal injury to the trachea and intratracheal instillation of benzo[a]pyrene. *Cancer Lett*, 29: 93–99. doi:10.1016/0304-3835(85)90128-4 PMID:4063958
- Rippe RM, Pott D (1989). *Kanzerogenitätsuntersuchungen von Nitro-PAH (Nitroarenen) im Hinblick auf ihre Bedeutung für die krebserzeugende Wirkung von Dieselmotorabgas*. In: *Gesellschaft zur Förderung der Luftthygiene und Silikoseforschung*. Düsseldorf: Stefan W. Albers, pp. 65–89.
- Rodriguez LV, Dunsford HA, Steinberg M *et al.* (1997). Carcinogenicity of benzo[a]pyrene and manufactured gas plant residues in infant mice. *Carcinogenesis*, 18: 127–135. doi:10.1093/carcin/18.1.127 PMID:9054599
- Rogan EG, RamaKrishna NVS, Higginbotham S *et al.* (1990). Identification and quantitation of 7-(benzo[a]pyren-6-yl)guanine in the urine and feces of rats treated with benzo[a]pyrene. *Chem Res Toxicol*, 3: 441–444. doi:10.1021/tx00017a009 PMID:2133095
- Roller M, Kamino K, Rosenbruch M (1992). *Carcinogenicity testing of bladder carcinogens and other organic compounds by the intraperitoneal and intravesicular route*. In: *Environmental Hygiene III*. Seemayer NH, Hadnagy W, editors. Berlin: Springer-Verlag, pp. 95–98.
- Ross JA, Nelson GB, Wilson KH *et al.* (1995). Adenomas induced by polycyclic aromatic hydrocarbons in strain A/J mouse lung correlate with time-integrated DNA adduct levels. *Cancer Res*, 55: 1039–1044. PMID:7866986
- Rossi L, Barbieri O, Sanguineti M *et al.* (1983). Carcinogenic activity of benzo[a]pyrene and some of its synthetic derivatives by direct injection into the mouse fetus. *Carcinogenesis*, 4: 153–156. doi:10.1093/carcin/4.2.153 PMID:6297822
- Ruggeri B, DiRado M, Zhang SY *et al.* (1993). Benzo[a]pyrene-induced murine skin tumors exhibit frequent and characteristic G to T mutations in the p53 gene. *Proc Natl Acad Sci U S A*, 90: 1013–1017. doi:10.1073/pnas.90.3.1013 PMID:8430068
- Saffiotti U, Montesano R, Sellakumar AR *et al.* (1972). Respiratory tract carcinogenesis in hamsters induced by different numbers of administrations of benzo[a]pyrene and ferric oxide. *Cancer Res*, 32: 1073–1081. PMID:4336025
- Sellakumar A, Stenbäck F, Rowland J (1976). Effects of different dusts on respiratory carcinogenesis in hamsters induced by benzo (a) pyrene and diethylnitrosamine. *Eur J Cancer*, 12: 313–319. doi:10.1016/0014-2964(76)90112-2 PMID:954792
- Sellakumar AR, Montesano R, Saffiotti U, Kaufman DG (1973). Hamster respiratory carcinogenesis induced by benzo[a]pyrene and different dose levels of ferric oxide. *J Natl Cancer Inst*, 50: 507–510. PMID:4702121
- Senft AP, Dalton TP, Nebert DW *et al.* (2002). Mitochondrial reactive oxygen production is dependent on the aromatic hydrocarbon receptor. *Free Radic Biol Med*, 33: 1268–1278. doi:10.1016/S0891-5849(02)01014-6 PMID:12398935
- Shen YM, Troxel AB, Vedantam S *et al.* (2006). Comparison of p53 mutations induced by PAH o-quinones with those caused by anti-benzo[a]pyrene diol epoxide in vitro: role of reactive oxygen and biological selection. *Chem Res Toxicol*, 19: 1441–1450. doi:10.1021/tx0601206 PMID:17112231
- Shimada T (2006). Xenobiotic-metabolizing enzymes involved in activation and detoxification of carcinogenic polycyclic aromatic hydrocarbons. *Drug Metab*

- Pharmacokinet*, 21: 257–276. doi:10.2133/dmpk.21.257 PMID:16946553
- Shimizu Y, Nakatsuru Y, Ichinose M *et al.* (2000). Benzo[a]pyrene carcinogenicity is lost in mice lacking the aryl hydrocarbon receptor. *Proceedings of the National Academy of Sciences of the United States of America*, 97: 779–782. doi:10.1073/pnas.97.2.779 PMID:10639156
- Simioli P, Lupi S, Gregorio P *et al.* (2004). Non-smoking coke oven workers show an occupational PAH exposure-related increase in urinary mutagens. *Mutat Res*, 562: 103–110. PMID:15279833
- Smith DM, Rogers AE, Herndon BJ, Newberne PM (1975a). Vitamin A (retinyl acetate) and benzo[a]pyrene-induced carcinogenesis in hamsters fed a commercial diet. *Cancer Res.*, 35: 11–16. PMID:162856
- Smith DM, Rogers AE, Newberne PM (1975b). Vitamin A and benzo[a]pyrene carcinogenesis in the respiratory tract of hamsters fed a semisynthetic diet. *Cancer Res*, 35: 1485–1488. PMID:1131819
- Solt DB, Polverini PJ, Calderon L (1987). Carcinogenic response of hamster buccal pouch epithelium to 4 polycyclic aromatic hydrocarbons. *Journal of Oral Pathology*, 16: 294–302. doi:10.1111/j.1600-0714.1987.tb00697.x PMID:2445943
- Sparnins VL, Mott AW, Barany G, Wattenberg LW (1986). Effects of allyl methyl trisulfide on glutathione S-transferase activity and BP-induced neoplasia in the mouse. *Nutrition and Cancer*, 8: 211–215. doi:10.1080/01635588609513895 PMID:3737423
- Stansbury KH, Flesher JW, Gupta RC (1994). Mechanism of aralkyl-DNA adduct formation from benzo[a]pyrene in vivo. *Chem Res Toxicol*, 7: 254–259. doi:10.1021/tx00038a019 PMID:8199315
- Steinhoff D, Mohr U, Hahneemann S (1991). Carcinogenesis studies with iron oxides. *Exp Pathol*, 43: 189–194. PMID:1797572
- Stenbäck F & Rowland J (1978). Role of particle size in the formation of respiratory tract tumors induced by benzo(a)pyrene. *Eur J Cancer*, 14: 321–326. doi:10.1016/0014-2964(78)90200-1 PMID:656185
- Stenbäck F & Rowland J (1979). Experimental respiratory carcinogenesis in hamsters: environmental, physico-chemical and biological aspects. *Oncology*, 36: 63–71. doi:10.1159/000225320 PMID:223099
- Stenbäck F, Sellakumar A, Shubik P (1975). Magnesium oxide as carrier dust in benzo(a)pyrene-induced lung carcinogenesis in Syrian hamsters. *J Natl Cancer Inst*, 54: 861–867. PMID:1127716
- Surh YJ, Liem A, Miller EC, Miller JA (1989). Metabolic activation of the carcinogen 6-hydroxymethylbenzo[a]pyrene: formation of an electrophilic sulfuric acid ester and benzylic DNA adducts in rat liver in vivo and in reactions in vitro. *Carcinogenesis*, 10: 1519–1528. doi:10.1093/carcin/10.8.1519 PMID:2752526
- Thaiparambil JT, Vadhanam MV, Srinivasan C *et al.* (2007). Time-dependent formation of 8-oxo-deoxyguanosine in the lungs of mice exposed to cigarette smoke. *Chem Res Toxicol*, 20: 1737–1740. doi:10.1021/tx700289g PMID:18031018
- Thyssen J, Althoff J, Kimmerle G, Mohr U (1981). Inhalation studies with benzo[a]pyrene in Syrian golden hamsters. *J Natl Cancer Inst*, 66: 575–577. PMID:6937711
- Toth B (1980). Tumorigenesis by benzo[a]pyrene administered intracolonicly. *Oncology*, 37: 77–82. doi:10.1159/000225408 PMID:7360483
- Urso P & Gengozian N (1984). Subnormal expression of cell-mediated and humoral immune responses in progeny disposed toward a high incidence of tumors after in utero exposure to benzo[a]pyrene. *J Toxicol Environ Health*, 14: 569–584. doi:10.1080/15287398409530606 PMID:6239929
- Van Duuren BL, Katz C, Goldschmidt BM (1973). Cocarcinogenic agents in tobacco carcinogenesis. *J Natl Cancer Inst*, 51: 703–705. PMID:4765384
- van Oostrom CT, Boeve M, van Den Berg J *et al.* (1999). Effect of heterozygous loss of p53 on benzo[a]pyrene-induced mutations and tumors in DNA repair-deficient XPA mice. *Environ Mol Mutagen*, 34: 124–130. doi:10.1002/(SICI)1098-2280(1999)34:2/3<124::AID-EM11>3.0.CO;2-F PMID:10529736
- Vesselinovitch SD, Kyriazis AP, Mihailovich N, Rao KVN (1975a). Factors influencing augmentation and/or acceleration of lymphoreticular tumors in mice by benzo[a]pyrene treatment. *Cancer Res*, 35: 1963–1969. PMID:1097103
- Vesselinovitch SD, Kyriazis AP, Mihailovich N, Rao KVN (1975b). Conditions modifying development of tumors in mice at various sites by benzo[a]pyrene. *Cancer Res*, 35: 2948–2953. PMID:1182688
- Vineis P & Husgafvel-Pursiainen K (2005). Air pollution and cancer: biomarker studies in human populations. *Carcinogenesis*, 26: 1846–1855. doi:10.1093/carcin/bgi216 PMID:16123121
- Von Tungeln LS, Xia Q, Herreno-Saenz D *et al.* (1999). Tumorigenicity of nitropolycyclic aromatic hydrocarbons in the neonatal B6C3F1 mouse bioassay and characterization of ras mutations in liver tumors from treated mice. *Cancer Letters*, 146: 1–7. doi:10.1016/S0304-3835(99)00206-2 PMID:10656603
- Warshawsky D & Barkley W (1987). Comparative carcinogenic potencies of 7H-dibenzo[c,g]carbazole, dibenz[a,j]acridine and benzo[a]pyrene in mouse skin. *Cancer Letters*, 37: 337–344. doi:10.1016/0304-3835(87)90119-4 PMID:3677065
- Warshawsky D, Barkley W, Bingham E (1993). Factors affecting carcinogenic potential of mixtures. *Fundamental and Applied Toxicology*, 20: 376–382. doi:10.1006/faat.1993.1048 PMID:8504912
- Watanabe KH, Djordjevic MV, Stellman SD *et al.* (2009). Incremental lifetime cancer risks computed for benzo[a]pyrene and two tobacco-specific N-nitrosamines in mainstream cigarette smoke compared with lung

- cancer risks derived from epidemiologic data. *Regul Toxicol Pharmacol*, 55: 123–133. PMID:19540296
- Wei SJ, Chang RL, Merkler KA *et al.* (1999). Dose-dependent mutation profile in the c-Ha-ras proto-oncogene of skin tumors in mice initiated with benzo[a]pyrene. *Carcinogenesis*, 20: 1689–1696. doi:10.1093/carcin/20.9.1689 PMID:10469612
- Wenzel-Hartung R, Brune H, Grimmer G *et al.* (1990). Evaluation of the carcinogenic potency of 4 environmental polycyclic aromatic compounds following intrapulmonary application in rats. *Exp Pathol*, 40: 221–227. PMID:1711479
- Weyand EH, Chen Y-C, Wu Y *et al.* (1995). Differences in the tumorigenic activity of a pure hydrocarbon and a complex mixture following ingestion: benzo[a]pyrene vs manufactured gas plant residue. *Chemical Research in Toxicology*, 8: 949–954. doi:10.1021/tx00049a008 PMID:8555410
- Wijnhoven SWP, Kool HJM, van Oostrom CTM *et al.* (2000). The relationship between benzo[a]pyrene-induced mutagenesis and carcinogenesis in repair-deficient Cockayne syndrome group B mice. *Cancer Res*, 60: 5681–5687. PMID:11059760
- Wislocki PG, Bagan ES, Lu AY *et al.* (1986). Tumorigenicity of nitrated derivatives of pyrene, benz[a]anthracene, chrysene and benzo[a]pyrene in the newborn mouse assay. *Carcinogenesis*, 7: 1317–1322. doi:10.1093/carcin/7.8.1317 PMID:3731386
- Wojdani A & Alfred LJ (1984). Alterations in cell-mediated immune functions induced in mouse splenic lymphocytes by polycyclic aromatic hydrocarbons. *Cancer Res*, 44: 942–945. PMID:6420057
- Xiao H, Rawal M, Hahm ER, Singh SV (2007). Benzo[a]pyrene-7,8-diol-9,10-epoxide causes caspase-mediated apoptosis in H460 human lung cancer cell line. *Cell Cycle*, 6: 2826–2834. PMID:17986867
- Xue W & Warshawsky D (2005). Metabolic activation of polycyclic and heterocyclic aromatic hydrocarbons and DNA damage: a review. *Toxicol Appl Pharmacol*, 206: 73–93. doi:10.1016/j.taap.2004.11.006 PMID:15963346
- Yu D, Kazanietz MG, Harvey RG, Penning TM (2002). Polycyclic aromatic hydrocarbon o-quinones inhibit the activity of the catalytic fragment of protein kinase C. *Biochemistry*, 41: 11888–11894. doi:10.1021/bi020270p PMID:12269833